



Hedmark University College

Faculty of applied ecology and agriculture

BRAGE

Hedmark University College's Open Research Archive

<http://brage.bibsys.no/hhe/>

This is the author's version of the article published in

Aquatic Ecology

The article has been peer-reviewed, but does not include the publisher's layout, page numbers and proof-corrections

Citation for the published paper:

Volk, C. & Kiffney, P. (2011). Comparison of fatty acids and elemental nutrients in periphyton, invertebrates, and cutthroat trout (*Oncorhynchus clarki*) in conifer and alder streams of western Washington state. *Aquatic ecology*, 46(1), 85-99.

DOI: 10.1007/s10452-011-9383-7

1 **Comparison of fatty acids and elemental nutrients in periphyton, invertebrates, and**
2 **cutthroat trout (*Oncorhynchus clarki*) in conifer and alder streams of western Washington**
3 **state**

4

5

6

7

8 Carol Volk¹

9 ¹South Fork Research, Inc.

10 44842 SE 145th St

11 North Bend, WA 98045

12 carol@southforkresearch.org

13

14 Peter Kiffney^{2,3}

15 ²NOAA-Fisheries, National Marine Fisheries Service, Northwest Fisheries Science Center, Fish

16 Ecology Division, Mukilteo Biological Field Station

17 10 Park Avenue, Building B

18 Mukilteo, WA 98275

19 Peter.kiffney@noaa.gov

20 and

21 ³ Hedmark University College, Department of Forestry and Wildlife Management, Campus

22 Evenstad, Elverum, Norway 2409

23

24 Keywords: fatty acids, trophic levels, nitrogen, phosphorus, alder, nutrients

25

Abstract

26
27 Organism growth and reproduction are often limited by nutrient availability in freshwater
28 ecosystems where, in some cases, food webs are primarily supported by allochthonous organic
29 matter. Therefore, we hypothesized that the composition of riparian vegetation would influence
30 the variability of N, P and fatty acid content of in-stream consumers. Specifically, we predicted
31 that organisms living in alder streams would have higher levels of N, P, and polyunsaturated
32 fatty acids than organisms in coniferous streams. To determine this, we sampled fresh and aged
33 leaf litter, periphyton, invertebrates, and cutthroat trout (*Oncorhynchus clarki*) from 6 streams in
34 western Washington state: 3 streams had high densities of nitrogen-fixing red alder (*Alnus rubra*)
35 in the riparian zone, whereas 3 had high densities of conifers. We found fresh alder litter had
36 twice the total polyunsaturated fatty acid concentrations of hemlock vegetation while there were
37 few statistical differences among aged alder and aged hemlock vegetation. Multidimensional
38 plots showed fatty acid profiles were unique to vegetation and fish while periphyton and
39 invertebrates shared the same multidimensional space. We used a mixed model to determine the
40 relative importance of vegetation type (fixed factor: conifer or alder), trophic levels (fixed factor:
41 periphyton, primary consumer, or fish) and streams (random factor) on individual fatty acid
42 concentrations. Total polyunsaturated fatty acids, 16:0, 20:1, 20:3n6 and total n3 were the only
43 fatty acids influenced by stream vegetation (vegetation + stream model or full model. 67% of
44 the fatty acids were best supported by the trophic +stream model. Nitrogen, P, Ca, Fe, C:N, N:P
45 and C:N:P were all best supported by the trophic level + stream model and Zn was the only
46 nutrient supported best by the full model. Correlations of n3 and n6 fatty acid concentrations
47 between periphyton and primary consumers, and primary consumers with trout indicated several
48 fatty acid metrics, such as n3:n6, showed food resources may affect relative fatty acid

49 abundances of consumers. Although vegetation type did not influence relative fatty acids of
50 stream organisms, the importance of trophic level likely indicates organisms have different
51 physical requirements for fatty acids. The significance of a random factor, 'stream,' suggests
52 that the relative abundances of fatty acids in periphyton, invertebrates and trout are more related
53 than similar organisms from another stream.

54

55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77

Introduction

There is a strong connection between riparian vegetation and forest stream food webs (Cummins et al. 1989; Richardson 1990; Kiffney et al. 2003). In general, food webs in these streams depend on biomolecules from terrestrial sources, such as leaf litter or soil runoff (Vannote et al. 1980; Barlocher 1992; Webster and Meyer 1997). The availability and quality of riparian leaf litter varies widely, and this variability potentially influences consumer populations (Volk 2004). Therefore, quantitatively assessing the chemical quality of a stream food web might be useful for predicting in-stream production and fish growth. Food quality is commonly assessed using C:N and N:P ratios, but essential fatty acids are an alternative measure of food quality that has recently been applied to lake and stream ecosystems (Arts 1998; Arts et al. 2009). Fatty acid indicators are unique in that many animals, including humans, lack the desaturation enzymes that act at the n3 and n6 positions of polyunsaturated fatty acids (PUFAs). These fatty acids are critical for hormone production and membrane fluidity, and since they cannot be produced they are essential dietary nutrients (Sargent et al. 1999). Furthermore, high dietary concentrations of PUFAs, specifically n3 and n6 fatty acids, promote growth and reproductive rates for aquatic invertebrates (Ravet et al. 2003; Brett et al. 2006). Few fatty acid studies have assessed fatty acid compositions across trophic levels in natural systems (Torres-Ruiz et al. 2007) or environmental factors that might affect fatty acid composition (e.g. Peeters et al. 2004) even though general fatty acid profiles of algae, invertebrates and fish are well summarized by Arts et al. (2009).

Streams in the Pacific Northwest are generally oligotrophic and, depending on underlying geology, can be limited by N, P or NP co-limited (Volk et al. 2008; Kiffney 2008; Sanderson et al. 2009). Red alder, *Alnus rubra*, is a common nitrogen-fixing species found along riparian

78 corridors and disturbed landscapes of the Pacific Northwest. Alder leaf litter and underlying
79 soils are rich in N and P, and our earlier research showed annual detrital inputs were about $3.5 \times$
80 higher in streams dominated by riparian red alder relative to streams bordered primarily by
81 conifers (Volk 2004). Concentrations of a number of important biomolecules (e.g., N, P) in
82 water and fluvial particulate organic matter were also higher in alder-dominated streams. Alder
83 additions occur throughout the year with a large pulse during leaf fall in autumn, resulting in
84 annual total inputs of N and P to select streams of the Olympic Peninsula, WA, USA of 8.0 and
85 $0.25 \text{ g/m}^2/\text{year}$, respectively (Volk 2004), which are 5 to $8 \times$ higher than inputs to nearby
86 conifer dominated streams. Others have also shown that riparian alder forests are associated with
87 variability in the trophic productivity of freshwater ecosystems in the western US (Goldman
88 1961; Compton et al. 2003; Volk et al. 2003).

89 Therefore, we hypothesized the chemistry of detrital subsidies from riparian vegetation
90 may affect the chemical composition of local aquatic biota since N, P and fatty acids are
91 essential for survival, growth, and reproduction (Müller-Navarra 1995; Bendiksen et al. 2003).
92 To test this hypothesis, we quantified concentrations of PUFAs, C, N, P, Ca, Fe, and Zn in fresh
93 vegetation, stream-aged leaf litter, periphyton, invertebrates, and coastal cutthroat trout
94 (*Oncorhynchus clarki*) from 6 independent watersheds in western Washington: three stream
95 riparian corridors were dominated by alder and three by coniferous vegetation. Our objectives
96 were to compare n3 and n6 fatty acid concentrations and elemental nutrient concentrations of: 1)
97 food webs in alder and conifer-dominated streams; and 2) among trophic levels.

98

99

Methods

100 *Study sites*

101 Six headwater tributaries to the Hoh (47° 48' 36", -124° 5' 12") and Clearwater rivers (47° 42' 7",
102 -124° 10' 23") on the western Olympic Peninsula in Washington state were used as study sites
103 (Table 1). We selected these sites because of the predominance of alder or conifer within 30 m
104 of the stream bank, low accessibility to anadromous salmon and vehicle accessibility. Bridge
105 Creek, Bull Creek and Hook Creek were classified as coniferous streams because their riparian
106 corridors were dominated by ~75 year old second-growth Sitka spruce (*Picea sitchensis*),
107 western redcedar (*Thuja plicata*), western hemlock (*Tsuga heterophylla*), and Douglas-fir
108 (*Pseudotsuga mensizeii*). Christmas, Shale, and Maple Creeks were classified as alder streams
109 and were predominantly vegetated with red alder within the riparian corridor. Classifications
110 were assessed by measuring litter flux to streams using 5 baskets placed within the bankfull
111 channel of a 200 m reach; alder sites were required to have over 90% of the litterfall composed
112 of alder vegetation. We calculated the catchment area upstream of sample sites (NHDPlus
113 hydrography) and then the percent of land within this catchment covered by hardwood species
114 (data from Landsat Vegetation Mapping (1998) and GAP vegetation coverages (1991)) (Table
115 1). Alder is the dominant hardwood within these coastal streams and we considered the
116 hardwood and broadleaf GIS layers a reasonable proxy for alder composition within watersheds.

117 *Sample collections*

118 In September 2003, alder and hemlock vegetation, periphyton, invertebrates, and
119 cutthroat trout were collected from a 200m reach of each stream for C, N, P, Fe, Ca, Zn and fatty
120 acid analyses. Freshly senesced alder leaves and hemlock needles were shaken from three trees
121 of each species and collected. Periphyton was scraped from rocks (5 samples/stream) with a
122 toothbrush, rinsed with deionized water, filtered onto Whatman GF/F) filters, and frozen.
123 Baetidae ($n_{\text{alder}} = 3$ individuals from a total of 2 streams, $n_{\text{conifer}} = 7$ individuals from a total of 2

124 streams), Heptageniidae ($n_{\text{alder}} = 11$ from 2 a total of 2 streams, $n_{\text{conifer}} = 4$ from 1 stream), and
125 Glossosomatidae ($n_{\text{alder}} = 6$ individuals from a total of 2 streams, $n_{\text{conifer}} = 6$ individuals from a
126 total of 2 streams) invertebrates were hand collected and frozen. Macroinvertebrate collections
127 were limited to these predominantly herbivorous and dominant families due to a limited
128 abundance of detritivores and shredders at the time of sampling. Five to ten cutthroat trout (fork
129 length of 6-15 cm) were collected from each stream using a single pass electrofishing survey,
130 weighed (nearest 0.1 g), measured (nearest 1 mm), and frozen. Single-pass electrofishing was
131 deemed a sufficient capture method as no fish population abundance estimates were planned for
132 the study (e.g. Bateman et al. 2005). All samples were frozen at -80°C until fatty acid extraction
133 and C, N and P and micronutrient analyses.

134 In addition to freshly senesced vegetation, we aged leaf litter in streams to simulate in-
135 stream detritus. We placed 10 g packets of senesced hemlock or alder in mesh bags (15 cm
136 diameter bags with 0.4 mm mesh) and zip-tied packets to large rocks in Christmas (alder
137 vegetation type) and Hook (conifer vegetation type) Creeks (note only one stream for each
138 vegetation type was used for packet placement and we considered individual packets as
139 replicates). Packets were submerged on 9/11/2003 and after 18 days packs were recovered and
140 placed in plastic bags for biomass measurements. We submerged leaf packs long enough to
141 allow diffusion of most nutrients and partial mass loss, as most leaf litter nutrients are lost within
142 24 hours of submersion in water (Gessner and Schwoerbel 1989) and 40% mass loss can be
143 found after four weeks (Braatne et al. 2007). In the lab, invertebrates were removed from packets
144 and remaining biomass was dried (30°C) to a constant weight and weighed. All weights used
145 within analyses are total biomass and were not corrected for inorganic matter accumulation.

146 *Fatty acid extraction*

147 A simultaneous extraction of wax esters and total fatty acids (Kattner and Fricke 1986;
148 Doerthe C. Müller-Navarra, University of Hamburg, Hamburg, Germany, personal
149 communication) was used for extraction of n3 and n6 fatty acids. Subsamples of leaf litter (fresh
150 and aged), periphyton, fish dorsal muscle tissue and whole invertebrates were freeze dried for 4
151 hours and weighed (sample weights ranged from 0.25-1.5mg, pending tissue type). Ten µl of the
152 internal standard, 21:0 (10mg/10ml methanol; Sigma #H-5149, heneicosanoic acid) was added
153 prior to three dichloromethane:methanol (2:1v/v) extractions. The first extraction was overnight
154 (15 hours) with 5ml of dichloromethane:methanol mixture, followed by a second and third
155 extraction of the sample material with 3 and 2ml of dichloromethane:methanol solution for 2-3
156 hours and 0.5 hours, respectively. Suspension liquid was removed, set aside, and chilled (32°C)
157 between extractions of source material. After all extractions were complete, set aside suspensions
158 were recombined and then evaporated with N₂ gas (30°C heat block), and resuspended with 2ml
159 of 3% sulfuric acid in methanol and 5ml 16% n-hexane addition. This mixture was heated for 4
160 hours at 80 °C, converting all Fatty Acids to Methyl Esters (FAMES), which are soluble in
161 hexane. FAMES were separated from the sulfuric acid matrix with four additional extractions
162 with n-hexane (water added to solution before extraction to dilute sulfuric acid matrix and
163 facilitate hexane:sulfuric acid solution separation), then N₂ gas evaporated to dryness and
164 dissolved in 1.5ml n-hexanes. All FAMES were frozen (-80 °C) until injection (5µl) into the gas
165 chromatograph (GC).

166 Fatty acids were analyzed on an HP6890 series GC with an Agilent DB-WAX (30 m) +
167 guard column (10m; 0.32 mm, 0.25 µm film) and PTV inlet. Gas chromatography program
168 specifics were 5 minutes at 40 °C (ramp rate = 10 °C/min), 5 minutes at 150 °C (ramp rate = 2
169 °C/min), 24 minutes at 220 °C (ramp rate = 2°C/min). The program was optimized for 18C fatty

170 acids and had a detection limit of 0.112mg dry weight. All program specifications were per the
171 University of California-Davis (Goldman Laboratory) specifications from Kattner and Fricke
172 (1986) and Doerthe C. Müller-Navarra (University of Hamburg, Hamburg, Germany, personal
173 communication). All fatty acids with retention times between 20 and 65 minutes were identified
174 by retention time on the chromatograph with comparison to reference standard (37 FAME,
175 Supelco Mix C4-C24). Reference peaks of interest included 10:0, 11:0, 12:0, 13:0, 14:0, 14:1,
176 15:0, 15:1, 16:0, 16:1, 17:0, 17:1, 18:0, 18:1n9, 18:2n6 cis and trans, 18:3n6, 18:3n3, 18:4n3,
177 20:0, 20:1, 20:2n*, 20:3n6, 20:4n6, 20:3n3 20:5n3 22:0, 22:1n9, 22:2n6, 23:0 24:0, 22:6n3, and
178 24:1. Once peaks were identified through comparison with the reference standard, areas for each
179 sample peak were corrected for the recovery volume of the internal standard, 21:0, and
180 multiplied by the total amount of sample (mg). No inferences on non-reference peaks were
181 made. Fatty acid analyses were replicated only within the study design (e.g. 6 detritus samples
182 per stream) and not for fatty acid analyses (e.g. 1 sample from stream extracted and run through
183 the GC multiple times). At least two blanks were included in each extraction and an additional 2
184 standards were included in each GC sample run.

185 *Elemental Nutrients*

186 Fresh alder and hemlock, aged alder and hemlock, periphyton, and trout muscle tissue
187 were analyzed for C, N, P, Ca, Fe, K, and Zn. Material for elemental nutrient analyses was
188 extracted from the same samples as fatty acids when there was ample material. Tissues were
189 freeze dried for 4 hours and ground for C and N analysis in a CE440 Elemental Analyzer
190 (Leeman Labs, Inc., University of Washington Oceanography Technical Services Laboratory,
191 Seattle, WA). To determine P, Ca, Fe, and Zn concentrations, freeze dried material was digested
192 with nitric acid for 12 hours, heated to 120 °C for 1 hour, oxidized with H₂O₂ until colorless and

193 resuspended in 5% HCl (modified P digest from Jones et al. (1991)). Samples were run on an
194 ICP (Inductively Coupled Plasma Analyzer, NOAA, Northwest Fisheries Science Center,
195 Seattle, WA) and nutrient concentrations were calculated using standard curves for laboratory
196 standards.

197 *Physical measurements*

198 One surface water grab sample in (September 2003) was collected for total N, total P,
199 ammonium, nitrite, nitrate, and phosphate analyses (Valderrama 1981). All water samples were
200 frozen (-80 °C) and analyzed within 1 month of field collection. Additional physical habitat
201 details from a 200m survey of each stream during August-September 2003 are summarized in
202 Table 1 and described in detail in Volk (2004).

203 *Statistical analyses*

204 Since we assumed individual plants and leaf packs were independent samples, a one-way
205 ANOVA was used to compare nutrient composition (fatty acids or elemental nutrients) between:
206 a) fresh alder and fresh conifer vegetation and b) aged alder and aged conifer detritus. All data
207 were tested for normality (Shapiro test and Q-Q plots) and non-normal data were $\ln(x)$
208 transformed to meet assumptions of normality. Low C:N ratios and high Ca, N, P, Fe, and Zn
209 content were used as indicators of food quality for comparisons within the study.

210 Fatty acid and elemental nutrient data for fresh alder and hemlock, aged alder and
211 hemlock, periphyton, Baetidae, Glossosomatidae, Heptageniidae, and trout were compared
212 among all six streams using three mixed models in an information-theoretical approach
213 (Burnham and Anderson 1998). With fatty acids and elemental nutrient data as response metrics,
214 the models were designed such that vegetation type (classified by alder or conifer-dominated
215 vegetation) and trophic level (fresh vegetation-alder or hemlock, aged vegetation-alder or

216 hemlock, periphyton, invertebrates and fish) were used as predictive, fixed factors. ‘Stream’ was
217 used as a random factor to capture inherent natural differences among streams (aka sites).

218 Models

219 Model A (full model):

220 Response = vegetation + trophic level + stream

$$221 \quad y_i = \alpha_{\text{veg } i} + \beta_{\text{trophic } i} + b_i + \epsilon_i$$

222 Model B: Response = vegetation + stream

$$223 \quad y_i = \alpha_{\text{veg } i} + b_i + \epsilon_i$$

224 Model C: Response = trophic level + stream

$$225 \quad y_i = \beta_{\text{trophic } i} + b_i + \epsilon_i$$

226

227 y_i is the i th response data point across all streams

228 $\alpha_{\text{veg } i}$ has two values: alder and conifer

229 $\beta_{\text{trophic } i}$ has five values: vegetation, detritus, periphyton, invertebrates, and trout

$$230 \quad b_i \sim N(0, \sigma_{\text{stream}})$$

$$231 \quad \epsilon_i \sim N(0, \sigma_{\text{residual}})$$

232

233 Response metrics were averages of each fatty acid for each trophic level in a stream and were

234 transformed to meet assumptions of normality prior to analyses. Most fatty acids were

235 transformed with log, square root, or cubed root transformations. We used arcsine-root

236 transformations for C, N and P percentages. Akaike’s Information Criteria corrected for small

237 sample sizes (AICc) were compared among models to assess the relative importance of

238 vegetation type and trophic level on nutrient composition of sampled organisms. The relative

239 Akaike weight (w_i) is the relative likelihood of each model divided by the sum of all weights for
240 all models and was calculated for each model. We considered models with greater than 0.70
241 relative weights as strongly supported (AICc differences >2 from best model), relative weights
242 between 0.40 and 0.70 as moderately supported (AIC differences between 1 and 2 from best
243 model), and models with less than 0.40 relative weights as minimally supported by the dataset
244 (Burnham and Anderson 1998). Because vegetation type was a binary fixed factor (alder or
245 conifer), support for this model indicated alder and conifer response metrics are different.
246 Coefficients of the model were used to determine directionality associated with alder and conifer
247 differences.

248 We utilized a multi-dimension scaling (MDS) plot of $\ln(x+1)$ to describe the relative
249 fatty acid composition of in-stream organisms in multivariate space (Primer 6 Software, Clarke
250 and Gorley 2006). To investigate whether fatty acids are correlated between trophic levels, we
251 created a correlation matrix of stream averages of periphyton ($n = 6$) and primary consumers
252 Heptageniidae ($n = 3$), Baetidae ($n = 3$), Glossosomatidae ($n = 4$) and trout ($n = 6$) for each fatty
253 acid. Since not all invertebrate families were found in all streams, n values varied among
254 invertebrate families. We also correlated all primary consumers with trout similarly to
255 periphyton and invertebrates. Pearson's correlation coefficients were used to determine
256 significance of correlations.

257

258

Results

259

260 *Comparison of fatty acids among trophic levels and vegetation types*

261 Of the three models, the trophic + stream model (Model C) best explained the variability
262 for 67% of the fatty acids (Table 2). All monosaturated fatty acids and 7 of 9 saturated fatty acids
263 were best supported by this model. The n3 and n6 fatty acids were of particular interest as they
264 are essential fatty acids; half of the n3 and 5 of 6 of the n6 fatty acids were strongly supported by
265 this model, all with relative weights greater than 0.70 (Table 2). We found similar results with
266 elemental nutrients, where 7 of 8 elemental nutrients or nutrient ratios (e.g. N:P) were best
267 supported by the trophic + stream model (Table 2).

268 Mean relative abundances of individual fatty acids were similar between alder and
269 conifer streams for periphyton, invertebrates and trout (Appendix 1). Although we expected
270 relative abundances of fatty acids would be influenced by riparian vegetation type, there was
271 almost no evidence that vegetation type influenced variation in fatty acid profiles, as the
272 vegetation + stream model (Model B) had low w_i values compared to the full and trophic +
273 stream models (Table 2). Two fatty acids were best supported by Model B (20:1 and 20:3n6)
274 and 3 fatty acid metrics (16:0, n3 and total PUFA) by the full model (Model A), but only five of
275 these showed relative abundances higher in alder than conifer; total PUFA was marginally higher
276 in conifer streams compared to alder streams. Despite limited support for relative abundances of
277 fatty acids, the influx of fatty acids from alder vegetation into streams may still be important to
278 stream ecosystems. We calculated the annual biomass inputs of fatty acids from senesced
279 vegetation by multiplying relative abundances of fatty acids by the estimated biomass of litterfall
280 from alder and conifer streams (Volk 2004) (Appendix 2). Annual alder biomass inputs were
281 ~3.5 times greater than conifer biomass inputs, this difference in detrital flux offers some
282 perspective on the total contribution of fatty acids to stream ecosystems from alder forests.

283 The general patterns of relative abundances of fatty acids among trophic levels can be
284 seen in Table 3. Invertebrates and trout had higher relative abundances of n3 fatty acids than
285 periphyton. Trout had highest levels of 22:6n3 relative to all other trophic levels. Vegetation and
286 periphyton had higher relative abundances of n6 fatty acids than invertebrates and trout. Relative
287 abundances of saturated fatty acids (SAFA) were also highest in vegetation and declined as
288 trophic level increased (Table 3).

289 There was some evidence that fatty acids were conserved in these stream food webs. The
290 n3:n6 ratio was positively correlated between periphyton and consumers ($r = 0.67, p = 0.03, n$
291 $= 10$, Figure 1) and consumers and trout ($r = 0.57, p = 0.08, n = 10$) (Figure 2). Furthermore, the
292 relative concentration of 18:3n3, was positively correlated between periphyton and consumers (r
293 $= 0.76, p < 0.01, n = 10$). Variability in 18:3n3 were positively correlated between consumers
294 and trout but this was not statistically significant ($r = 0.53, p = 0.11, n = 10$). Percent PUFA was
295 also positively correlated between primary consumers and trout ($r = 0.63, p = 0.03, n = 10$) but
296 not between periphyton and consumers ($r = 0.13, p = 0.70, n = 11$). The other n3 and n6
297 polyunsaturated fatty acids exhibited no correlations between different trophic levels (results not
298 shown).

299 Results from our multidimensional plots showed fatty acid profiles of trout were tightly
300 grouped and distinct from vegetation, periphyton and invertebrates. All vegetation (fresh alder,
301 aged alder, fresh hemlock and aged hemlock) showed considerable overlap, but were distinct
302 from periphyton, invertebrates and trout (Figure 3). The invertebrate families (Baetidae,
303 Heptageniidae, and Glossosomatidae) did not separate in multidimensional space. Similarly,
304 periphyton fatty acid profiles were highly variable and overlapped with all three invertebrate
305 families.

306

307 *Nutrient concentrations of decomposing alder and conifer litter in streams*

308 We compared fatty acid concentrations from the two streams (1 alder and 1 conifer) after
309 18 days of decomposition (samples were used as replicates). Eighteen carbon fatty acids were
310 10-30% lower in both aged alder and hemlock compared to fresh litter concentrations, and the
311 relative abundances of 50% of measured PUFAs dropped (Table 3). N3 and n6 fatty acid
312 concentrations decreased by 30% in alder but increased by 5-25% in hemlock (Figure 4 and
313 Table 3). In alder, 18:4n3 concentrations declined by 77%, but increased by 40% in hemlock,
314 while 18:2n6 concentrations were reduced in both hemlock and alder (44 and 50%, respectively)
315 (Figure 1). 18:3n3 was the only polyunsaturated fatty acid where aged alder concentrations were
316 statistically higher than aged hemlock concentrations ($n = 12, p < 0.01$).

317 Elemental concentrations of Ca, N and P were significantly higher in fresh alder than
318 hemlock vegetation, and C:N ratios were significantly lower (percent by weight) (Table 4).
319 Nitrogen, C, and Zn were significantly higher, and C:N ratios were almost 3 times lower in aged
320 alder detritus relative to aged hemlock. Percent P was not statistically different between aged
321 vegetation types ($n = 18, p = 0.75$, Table 4). In alder, 36-67% of Ca was lost over 18 days
322 whereas C loss rates were twice as fast in hemlock than alder (1.00 and 0.53% per day,
323 respectively).

324

325

Discussion

326 This is one of the first studies to investigate fatty acids in natural stream food webs, and
327 factors that may affect the variability of these essential biomolecules. Overall, most of the
328 variability in fatty acids was attributed to trophic level and stream, with vegetation type as an

329 important covariate for only a few fatty acids. We speculate the importance of trophic level and
330 stream suggests organisms were not limited in essential fatty acids and that metabolic differences
331 likely accounted for the significance of trophic level for almost all fatty acid and nutrient
332 response metrics.

333 One of the more intriguing results was the positive correlation among the relative
334 abundance of fatty acids in different trophic levels (periphyton, consumers, and fish). The
335 positive correlation between stream trophic levels for some fatty acids suggests consumers
336 reflect the chemical composition of their food resources. Specifically, levels of n3 and n6 fatty
337 acids in periphyton were positively correlated with fatty acids in primary consumers and trout,
338 suggesting these materials were conserved as they moved up the food chain. Feeding trials of
339 phytoplankton (cryptophytes, chlorophytes and cyanophytes) to *Daphnia pulex* showed high
340 correlations of 20:5n3 + 22:6n3 and 20:4n6 between food sources and *Daphnia* (Brett et al.
341 2006). Similarly, feeding trials using 18:2n6, 18:3n3, 20:4n6 or an n3 PUFA mix for laboratory
342 reared Arctic charr found a general dominance of n3 PUFA in fish muscle tissue when diets were
343 composed of 18:3n3 (Olsen et al. 1991), supporting our observations that a high PUFA content
344 in invertebrates correlated to high PUFA content in fish tissue. However, our study was
345 observational and exploratory, which limits our inference, but suggest some potential future
346 studies evaluating how fatty acids may influence food webs in stream ecosystems. For example,
347 does natural variation in essential fatty acids influence growth rate of stream fish through higher
348 growth efficiencies?

349 For elemental nutrients, lower C:N ratios in periphyton than terrestrial vegetation
350 indicates that aquatic primary producers were a higher quality food for consumers and predators
351 than terrestrial producers, and similar conclusions have been drawn by Reiners (1986) and Elser

352 et al. (2000). This is likely due to the high proportion of structural carbon material in terrestrial
353 vegetation (McGroddy et al. 2004). Furthermore, all else being equal, the lower C:N of algae
354 and alder may result in differences in growth efficiencies of higher trophic levels relative to
355 systems dependent on more recalcitrant sources of energy such as conifer needles, as conifer
356 needles break down more slowly than deciduous litterfall (Alberíño and Balseiro 2002). In both
357 terrestrial and lake environments, a reduction in the conversion efficiency of carbon into new
358 biomass ('gross growth efficiency') has been correlated with higher food C:N and C:P ratios
359 (Elser et al. 2000).

360 Trout, vegetation and invertebrates separated well in multidimensional space while
361 periphyton tended to overlap with vegetation and invertebrates. The high percentages of PUFA
362 (56%), and 22:6n3 (5 times higher relative abundance than other trophic levels) likely
363 contributed to the separation of trout from other trophic levels. Metabolic differences among
364 trout and invertebrates likely account for the separation of the two consumers, as the essential
365 fatty acid requirements for growth and cell structure are different (Ackman 1998). The
366 considerable overlap between periphyton and primary consumers was expected, as these
367 invertebrates were predominantly grazers, scrapers, and collectors that feed on periphyton
368 (Cummins 1973). In general, Baetidae feed by scraping algae and fine detritus from submerged
369 rocks or other submerged materials, such as woody debris. Heptageniidae are surface feeding
370 collectors or scrapers (mineral or organic scrapers) and Glossosomatidae are typically mineral
371 scrapers (Cummins 1973). The overlap between the fatty acid profiles of these invertebrate
372 families and periphyton is therefore not surprising and we assume periphyton is the dominant
373 food source for the sampled invertebrates in the study streams.

374 Results from this study suggest the relative abundance of a few fatty acids or n3:n6 ratios
375 of fatty acids can be used to assess the potential biological (e.g., growth, survival, productivity)
376 importance of different chemical components in stream food web (riparian plants, periphyton,
377 primary consumers, and trout. This suggests that streams with high relative abundance of these
378 fatty acids in periphyton will have similarly high abundances of these fatty acids in grazers and
379 trout. However, the majority of correlations among trophic levels for n3 and n6 fatty acids were
380 not significant, suggesting that some fatty acids may be better tracers than others. Further
381 studies considering feeding habits and trials in mesocosm environments with natural food
382 sources are needed especially those that quantify whether these fatty acids actually contribute to
383 variation in performance in higher trophic levels.

384 One of our main objectives of this study was to determine if leaf litter inputs from alder
385 vegetation influence the relative abundance of fatty acids of in-stream organisms because
386 primary producers are the only source of these fatty acids for higher trophic levels. We did find
387 higher abundances of PUFA, 18:3n3, 20:5n3, 20:4n6 and 22:6n3 fatty acids, and higher N and P
388 in leaf material from alder relative to hemlock vegetation (N and P results similar to Volk 2004).
389 However, the results presented here provided little evidence that fatty acids and elemental
390 nutrients in periphyton, invertebrates and trout were strongly influenced by vegetation type. We
391 suggest three potential reasons for this result: 1) the relative abundance of only 5 fatty acids was
392 higher in alder than conifer vegetation, suggesting there are very few fatty acids where we would
393 have expected to see influences from alder vegetation; 2) low sample sizes, especially for
394 invertebrates, did not provide enough power of detection for this study; and 3) polyunsaturated
395 fatty acids are not limiting food webs in these small streams and therefore relative abundances do
396 not change with the presence of additional fatty acid resources.

397 Instead, our model results indicated trophic level and streams were important covariates
398 predicting variation in biomolecules. The importance of trophic level likely indicates organisms
399 at different trophic levels have different metabolic requirements for fatty acids, particularly
400 essential fatty acids. The significance of a random factor, 'stream,' suggests that the relative
401 abundances of fatty acids in periphyton, invertebrates and trout are stream-specific and are
402 responding to local environmental or communal variables. Physical aspects or food resources
403 unique to each stream could influence the fatty acid profiles of these food webs. For example,
404 this 'stream effect' may be linked to differences in the composition of in-stream primary
405 producers, litterfall, other inputs of biomolecules (e.g. plant reproductive structures) or feeding
406 relationships. However we investigated a limited number of food resources and physical aspects
407 of a site that may contribute to this random variation in fatty acid profiles.

408 Aging leaf litter in streams reduced the relative abundances of n3 and n6 fatty acids, and
409 this may be due to an accumulation of inorganic material on aged leaf litter. This hypothesis may
410 be supported by the large increase in Fe between fresh and aged material, as Fe is common
411 element in inorganic minerals. Other changes in n3 and n6 fatty acid content between fresh and
412 aged material may have been due to algal or microbial colonization, but these communities were
413 not directly studied.

414

415

Conclusions

416 Relationships between 1) periphyton and invertebrate primary consumers and 2)
417 invertebrate primary consumers and fish suggest that the relative abundance of these resources
418 were conserved. Further observational and experimental studies are needed to improve our
419 understanding of the nutritional ecology of freshwater ecosystems, because this understanding

420 may help us conserve and restore ecologically and economically important fish species and their
421 ecosystems. Although the widespread abundance of riparian red alder in the Pacific Northwest
422 provides particulate and dissolved nutrient resources, it is difficult to discern the role of these
423 resources for primary and secondary consumers unless the limiting resources of the local
424 ecosystems are known. Moreover, we need a better understanding of the linkages between
425 watershed and riparian conditions that may affect chemical constituents potentially important in
426 the trophic productivity of freshwater food webs because this understanding may improve our
427 restoration and management of forested watersheds with economically and ecologically
428 important fish (Wipfli and Baxter 2010).

429

430

Acknowledgements

431 Significant assistance was provided by Anne Liston and the University of California Limnology
432 Laboratory, Michael Brett (University of Washington), and Brice Semmens (Scripps Institute of
433 Oceanography, University of California). Funding was provided by the National Oceanic and
434 Atmospheric Administration (Northwest Fisheries Science Center) and the U.S. Forest Service.

435

436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458

Literature Cited

Ackman RG (1998) Comparison of lipids in marine and freshwater organisms. In: Arts MT, Wainman BC (eds) *Lipids in Freshwater Ecosystems*, Springer, New York, pp 263-288

Alberíño RJ, Balseiro EG (2002) Leaf litter breakdown in Patagonian streams: native versus exotic trees and the effect of invertebrate size. *Aquat Conserv: Mar Freshwat Ecosyst* 12: 181-192

Arts MT (1998) Lipids in freshwater zooplankton: selected ecological and physiological aspects. In: Arts MT, Wainman BC (eds) *Lipids in Freshwater Ecosystems*, Springer, New York, pp 71-84

Arts MT, Brett MT, Kainz MJ (2009) *Lipids in aquatic ecosystems*. Springer, New York

Barlocher F (1992) Community organization. In: Barlocher F (ed) *The ecology of aquatic hyphomycetes*, Springer-Verlag, Berlin, pp 38-76

Bateman DS, Gresswell RE, Torgersen CE (2005) Devaluating single-pass catch as a tool for identifying spatial patterns in fish distribution: *J Fresh Ecology* 20:335-345

Bendicksen EÅ, Arnesen AM, Jobling M. (2003) Effects of dietary fatty acid profile and fat content on smolting and seawater performance in Atlantic salmon (*Salmo salar* L.). *Aquaculture* 225:149-163

459

460 Braatne JH, Sullivan SMP, Chamberlain E (2007) Leaf decomposition and stream
461 macroinvertebrate colonization of Japanese Knotweed, an invasive plant species. Internat Rev
462 Hydrobiol 92:656-665

463

464 Brett MT, Müller-Navarra DC, Ballantyne AP, Ravet JL, Goldman CR (2006) *Daphnia* fatty
465 acid content reflects that of their diet. Limnol Oceanogr 51:2428-2437

466

467 Burnham KP, Anderson DR (1998) Model selection and multimodel inference. Springer, New
468 York

469

470 Clarke KR, Gorley RN (2006) PRIMER v6: User Manual/Tutorial. PRIMER-E, Plymouth

471

472 Compton JE, Robbins Church M, Larned ST, Hogsett WE (2003) Nitrogen export from forested
473 watersheds in the Oregon Coast Range: the role of N₂-fixing red alder. Ecosystems 6:773-785

474

475 Cummins KW (1973) Trophic relations of aquatic insects. Ann Rev Entomol18: 183-206

476

477 Cummins KW, Wilzbach MA, Gates DM, Perry JB, Taliaferro WB (1989) Shredders and
478 riparian vegetation. BioScience 39:24-30

479

480 Elser JJ, Fagan WF, Denno RF, Dobberfuhl DR, Folarin A, Huberty A, Interlandi S, Kilham SS,
481 McCauley E, Schulz KL, Siemann EH, Sterner RW (2000) Nutritional constraints in terrestrial
482 and freshwater food webs. *Nature* 408:578-580
483
484 Gessner MO, Schwoerbel J (1989) Leaching kinetics of fresh leaf-litter with implications for the
485 current concept of leaf-processing in streams. *Archiv Hydrobiol* 115:81-90
486
487 Goldman CR (1961) The contribution of alder trees (*Alnus tenuifolia*) to the primary
488 productivity of Castle Lake, California. *Ecology* 42:282-288
489
490 Jones JB, Wolf B, Mills HA (1991) Plant analysis handbook: a practical sampling, preparation,
491 analysis, and interpretation guide. Micro-Macro Publishers, Athens, GA
492
493 Kattner G, Fricke HSG (1986). Simultaneous determination of fatty acids and alcohols in wax
494 esters of marine organisms. *J Chromat* 361:263-268
495
496 Kiffney PM, Richardson JS, Bull JP (2003) Responses of periphyton and insects to experimental
497 manipulation of riparian buffer width along forest streams. *J Appl Ecol* 40:1060-1076
498
499 Kiffney, PM (2008). Response of lotic producer and consumer trophic levels in headwater
500 streams of southwestern British Columbia in response to gradients of resource supply and
501 predation pressure. *Oikos* 114:1428-1440
502

503 McGroddy ME, Daufresne T, Hedin LO (2004) Scaling of C:N:P stoichiometry in forests
504 worldwide: implications of terrestrial Redfield-type ratios. *Ecology* 85:2390-2401
505

506 Müller-Navarra D (1995) Evidence that a highly unsaturated fatty acid limits *Daphnia* growth in
507 nature. *Archiv für Hydrobiol.* 132:297-307
508

509 Olsen RE, Henderson RJ, and Ringo E (1991) Lipids of Arctic charr, *Salvelinus alpinus* (L.) I.
510 Dietary induced changes in lipid class and fatty acid composition. *Fish Physiol Biochem* 9:151-
511 164
512

513 Peeters ETHM, Gylstra R, Vos JH (2004) Benthic macroinvertebrate community structure in
514 relation to food and environmental variables. *Hydrobiologia* 519:103-115
515

516 Ravet JL, Brett MT, Müller-Navarra DC (2003) A test of the role of polyunsaturated fatty acids
517 in phytoplankton food quality for *Daphnia* using liposome supplementation. *Limnol Oceanogr*
518 48:1938-1947
519

520 Reiners WA (1986) Complementary models for ecosystems. *Am Nat* 127:59-73
521

522 Richardson WB (1990) A Comparison of detritus processing between permanent and intermittent
523 headwater streams. *J Freshwat Ecol* 5: 341-357
524

525 Sanderson BL, Coe HJ, Tran CD, Macneale KH, Harstad DL, Goodwin AB (2009) Nutrient
526 limitation of periphyton in Idaho streams: results from nutrient diffusing substrate experiments. J
527 N Am Benthol Soc 28:832-845
528
529 Sargent J, Bell G, McEvoy L, Tocher D, Estevez A (1999) Recent developments in the essential
530 fatty acid nutrition of fish. Aquaculture 177:191-199
531
532 Torres-Ruiz M, Wehr JD, Perrone AA (2007) Trophic relations in a stream food web:
533 importance of fatty acid for macroinvertebrate consumers. J N Am Benthol Soc 26: 509-522
534
535 Valderrama JC (1981) The simultaneous analysis of total nitrogen and total phosphorus on
536 natural waters. Mar Chem 10:109-122
537
538 Vannote RL, Minshall G, Cummins KW, Sedell JR, Cushing CE (1980) The river continuum
539 concept. Can J Fish Aquat Sci 37:130-137
540
541 Volk CJ, Kiffney PM, Edmonds RL (2003) Role of riparian red alder (*Alnus rubra*) in the
542 nutrient dynamics of coastal streams of the Olympic Peninsula, WA.U.S.A. In Stockner JG (ed)
543 Nutrients in salmon ecosystems: sustaining production and biodiversity, American Fisheries
544 Society, Bethesda, pp 213-225
545
546 Volk CJ (2004) Nutrient and biological responses to red alder (*Alnus rubra*) presence along
547 headwater streams: Olympic Peninsula, WA. Dissertation, University of Washington

548

549 Volk CJ, Kiffney PM, Edmonds RL (2008) Nutrient limitation in red alder and conifer

550 forested streams of western Washington. *Am Mid Nat* 159:190-199

551

552 Webster JR, Meyer JL (1997) Stream organic matter budgets. *J N Am Benthol Soc* 16:3-161

553

554 Wipfli MS, Baxter CV (2010) Linking ecosystems, food webs, and fish production: Subsidies in

555 salmonid watersheds. *Fisheries* 35: 373-387

556 Table 1. Physical stream habitat characteristics of study sites

557

	Stream					
Vegetation	Maple Alder	Shale Alder	Christmas Alder	Bridge Conifer	Hook Conifer	Bull Conifer
Alder in watershed, %	16.0	25.0	62.0	46.0	0	0
Alder leaf litter in riparian area, %	100.0	97.0	91.0	10.0	34.0	18.0
Alder leaf litterfall in riparian area, g/m ²	15.2	59.8	81.5	1.8	12.5	8.2
Conifer leaf litterfall in riparian area, g/m ²	0	2.5	6.9	7.9	16.4	12.0
Stream gradient, %	2.7	2.4	1.2	5.1	3.6	1.8
Discharge, L/s	16.9	21.6	17.4	1.3	153.0	1.9
7d average T, °C	12.0	11.5	12.0	11.8	11.7	11.5
TP, µg/L	26.0	35.0	25.0	25.0	29.0	28.0
TN, µg/L	102.0	169.0	85.0	219.0	161.0	165.0
PO ₄ , µg/L	4.4	4.8	2.9	7.5	4.0	2.6
NO ₃ , µg/L	64.0	30.0	13.0	195.0	84.0	81.0
NH ₄ , µg/L	6.7	5.7	8.0	4.4	1.4	5.2

558

559 Table 2. AIC scores and relative model weights (w_i) for the full, vegetation and trophic models.
560 NA = Data not analyzed using mixed models due to high zero counts in data that led to violations
561 in normality assumptions. Response metrics are relative fatty acid abundance or elemental
562 nutrient concentrations per sample. * values calculated using subset of data for model due to
563 abundance of zeros within trophic levels. + data did not conform well to assumptions of
564 normality. ++ 3 models not completed due to limited amount of non-zero data in all trophic levels.
565

Response metric	AICc scores			w_i		
	Full model (Model A)	Vegetation + stream (Model B)	Trophic + stream (Model C)	Full model	Vegetation + stream	Trophic + stream
14:0	-26.29	28.49	-31.49	0.07	0.00	0.93
15:0	-181.63	-178.34	-187.56	0.05	0.01	0.94
16:0	756.96	799.82	757.76	0.60	0.00	0.40
17:0	-194.43	-172.64	-201.57	0.03	0.00	0.97
18:0	-47.20	-70.42	-52.86	0.00	1.00	0.00
20:0	133.15	165.79	129.53	0.14	0.00	0.86
22:0	191.45	207.96	187.86	0.14	0.00	0.86
23:0	NA	NA	NA	NA	NA	NA
24:0*	112.65	127.95	109.29	0.16	0.00	0.84
SAFA	-218.74	-173.69	-225.29	0.04	0.00	0.96
14:1 ⁺⁺	NA	NA	NA	NA	NA	NA
15:1 ⁺⁺	NA	NA	NA	NA	NA	NA
16:1	707.06	788.95	708.77			
17:1 ⁺	164.85	194.02	161.01	0.13	0.00	0.87
18:1n9	256.17	263.12	252.70	0.15	0.00	0.85
20:1	0.048	-6.60	-4.44	0.03	0.73	0.25
22:1n9 ⁺⁺	NA	NA	NA	NA	NA	NA
24:1 ⁺⁺	NA	NA	NA	NA	NA	NA
MUFA	-122.45	-88.43	-129.29	0.03	0.00	0.97
18:3 n 3	305.95	350.38	304.45	0.32	0.00	0.68
18:4 n 3	-40.38	-44.40	-45.83	0.04	0.32	0.64
20:5 n 3	43.93	170.12	38.62	0.07	0.00	0.93
20:3 n 3*	60.82	62.77	57.21	0.13	0.05	0.82
22:6 n 3 ⁺	203.65	469.65	201.25	0.23	0.00	0.77
n3	848.63	1009.01	850.91	0.76	0.00	0.24
18:2 n 6	-106.36	-47.67	-111.17	0.08	0.00	0.92
18:3 n 6	117.40	125.15	113.06	0.10	0.00	0.90
20:3 n 6 ⁺	68.43	63.51	65.085	0.06	0.65	0.30
20:4 n 6	-65.06	23.04	-69.87	0.08	0.00	0.92
22:2 n 6*	NA	51.90	NA	NA	NA	NA
n6	-97.02	-92.27	-102.92	0.05	0.00	0.95
n3:n6	-85.82	14.96	-92.06	0.04	0.00	0.96
PUFA	871.74	989.69	874.67	0.81	0.00	0.19
C ⁺	-153.33	-74.43	-160.53	0.03	0.00	0.97
N	-244.45	-96.48	-252.08	0.02	0.00	0.98
P	64.00	119.59	60.82	0.17	0.00	0.83
Ca	70.23	91.02	67.35	0.19	0.00	0.81
Fe	86.26	218.36	82.42	0.13	0.00	0.87
Zn	110.21	114.95	110.25	0.48	0.04	0.47
C:N	-188.55	-76.31	-193.77	0.07	0.00	0.93
N:P	36.50	46.72	34.48	0.27	0.00	0.73
C:N:P	38.25	123.95	32.74	0.06	0.00	0.94

569 Table 3. Summary of fatty acids in aquatic food web. Values are relative percentages of fatty
570 acid (or fatty acid ratio) averaged across all streams. N indicates the number of replicates (total
571 for all streams). Baetidae not collected from Hook Creek and Glossomatidae not collected from
572 Christmas Creek.

Fatty acid	Vegetation				Periphyton	Baetidae	Hept	Gloss	Trout
	Fresh Alder	Fresh hemlock	Aged alder	Aged hemlock					
n =	6	6	6	6	27	10	15	12	36
14:0	3.00	3.75	3.92	4.59	4.52	1.25	0.92	3.29	0.80
15:0	0.38	0.66	0.49	0.41	0.78	0.26	0.69	0.33	0.18
16:0	25.88	20.10	24.45	22.37	30.25	27.66	20.25	20.96	21.29
17:0	0.56	0.85	0.69	0.55	0.49	1.19	1.53	0.84	0.63
18:0	6.88	11.98	5.61	8.74	8.06	9.32	13.14	4.32	7.03
20:0	3.69	4.08	3.80	3.72	1.08	0.49	1.88	0.45	0.20
22:0	6.53	2.75	2.53	4.02	1.02	0.39	2.28	0.43	0.15
23:0	0.43	3.02	0.42	0.15	0.02	0	0	0	0
24:0	3.73	3.97	3.99	7.49	1.44	0.01	0.06	0	0.01
SAFA	51.09	51.17	45.89	52.05	47.67	40.59	40.74	30.63	30.29
14:1	0.35	0.32	0.07	0.16	0.02	0	0.01	0.40	0
15:1	0	0.81	0	0	0	0	0.01	0.02	0
16:1	1.48	2.38	2.13	1.33	11.9	5.98	11.23	10.53	3.21
17:1	0.13	0.84	0.33	0	1.32	0.30	0.31	1.67	0.08
18:1n9	12.68	14.76	14.10	14.87	10.53	8.49	10.21	10.49	9.58
20:1	0.30	0	0	0	0.12	0.05	0.01	0.06	0.08
22:1n9	0	0.26	0	0.49	0.02	0.05	0	0	0.01
24:1	0	0	0	0	0.19	0	0	0.20	0.02
MUFA	14.63	19.37	16.63	16.85	24.11	14.88	21.78	23.01	12.98
18:3 n 3	18.98	8.79	21.17	10.37	8.01	19.18	8.26	17.37	7.41
18:4 n 3	1.78	0.74	0.67	0.64	1.38	2.21	0.98	3.64	1.21
20:5 n 3	0.78	0	1.14	3.43	7.16	14.08	19.39	16.62	11.87
20:3 n 3	0.22	0.29	0.21	0	0.06	0.18	0.35	0.23	0.70
22:6 n 3	0.27	5.38	0	3.38	1.07	0.29	0.22	0.23	28.41
n3	22.03	15.20	23.19	17.82	17.67	35.95	29.20	38.09	49.61
18:2 n 6	10.90	11.20	7.71	8.36	6.21	6.85	3.99	3.83	3.99
18:3 n 6	0.09	0.32	0	0	0.42	0.23	0.19	0.52	0.07
20:3 n 6	0	0	0	0.38	0.02	0.07	0.01	0.34	0.25
20:4 n 6	0.44	0	0.50	0	1.69	14.08	3.62	3.51	2.66
22:2 n 6	0	0	3.46	4.25	1.26	0	0	0	0.06
n6	11.43	11.52	11.67	12.99	9.70	8.34	7.81	8.20	7.03
n3:n6	2.09	1.52	2.32	1.73	2.26	4.57	3.96	5.71	7.51
PUFA	34.20	28.92	37.21	30.82	27.94	44.53	37.30	46.30	56.71
SAFA: (MUFA+PUFA)	1.05	1.06	0.85	0.92	0.95	0.78	0.61	0.44	0.43

573

574

575 Table 4. Summary of elemental nutrient concentrations for fresh and aged alder and hemlock
 576 litter. Significance values are for ANOVA analyses comparing fresh alder to fresh hemlock and
 577 aged alder to aged hemlock ($\alpha = 0.05$). Fe and Zn values not available for fresh hemlock due to
 578 sample limitations.

579

	n	Fresh			Aged		
		Alder	Hemlock	p	Alder	Hemlock	p
C (%)	12	47.00	59.08	0.20	36.35	41.60	0.30
N (%)	12	2.37	0.99	0.02	3.06	1.41	0.03
P (%)	18	0.27	0.13	<0.01	0.32	0.22	0.75
C:N (molar)	12	23.13	69.61	<0.01	14.38	34.28	<0.01
C:P (molar)	12	459.81	1344.46	<0.01	6.03	23.89	0.04
C:N:P (molar)	12	2.7x10 ⁶	1.9x10 ⁶	<0.01	2266.05	14261.83	0.04
Ca ($\mu\text{g/g}$)	18	22846.00	4704.00	0.02	15302.00	5817.00	<0.01
Fe ($\mu\text{g/g}$)	18	322.23	<i>na</i>	<i>na</i>	3957.00	2333.00	0.21
Zn ($\mu\text{g/g}$)	18	252.08	<i>na</i>	<i>na</i>	4976.00	148.00	0.03

580

581

582 **Fig 1** Correlations between periphyton and primary consumers for a) 18:3n3; b)n6 and c) n3:n6
583 fatty acids

584

585 **Fig 2** Correlation between primary consumers and trout for n3:n6 fatty acids

586




587 **Fig 3** Fresh and aged leaf litter, periphyton, invertebrates and trout fatty acids in
588 multidimensional space


589

590 **Fig 4** Major fatty relative abundances in fresh and aged vegetation. n= 6 for each series. sum n6
591 = sum of 18:2n6 (cis and trans) 18:3n6, 20:3n6, 20:4n6, and 22:2n6; SAFA = saturated fatty
592 acids; MUFA= monosaturated fatty acids, PUFA = polyunsaturated fatty acids. Bars indicate
593 standard error, * indicates significance between fresh alder and fresh hemlock at $p < 0.05$ and +
594 indicates significance between aged alder and aged hemlock at $p < 0.05$


595

596






598
599



600



603 Appendix 1. Fatty acid summary for alder and conifer streams. Values are relative percentages of
 604 fatty acid (or fatty acid ratio) averaged across all streams. N indicates the number of replicates
 605 (total for all streams). Baetidae not collected from Hook Creek and Glossosomatidae not
 606 collected from Christmas Creek.
 607

Fatty acid	Periphyton Alder	Periphyton Conifer	Baetidae Alder	Baetidae Conifer	Hept Alder	Hept Conifer	Gloss Alder	Gloss Conifer	Trout Alder	Trout Conifer
n=	14	13	3	7	11	4	6	6	13	23
14:0	5.82	3.13	1.59	1.21	0.92	0.92	3.68	2.90	0.77	0.83
15:0	0.97	0.57	0.23	0.27	0.79	0.40	0.29	0.37	0.17	0.19
16:0	31.47	28.94	26.84	27.74	19.97	21.02	22.30	19.63	21.64	21.01
17:0	0.56	0.41	0.90	1.22	1.61	1.31	0.70	0.99	0.57	0.67
18:0	8.93	7.13	6.22	9.64	13.68	11.65	3.78	4.87	6.63	7.35
20:0	1.38	0.75	0.28	0.51	1.96	1.65	0.37	0.53	0.15	0.25
22:0	1.23	0.79	0.15	0.42	2.40	1.96	0.39	0.47	0.10	0.19
23:0	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
24:0	1.45	1.44	0.00	0.02	0.08	0.00	0.00	0.00	0.03	0.00
SAFA	51.86	43.16	36.21	41.02	41.41	38.91	31.50	29.75	30.05	30.50
14:1	0.04	0.00	0.00	0.00	0.01	0.00	0.06	0.02	0.00	0.00
15:1	0.00	0.00	0.00	0.00	0.01	0.00	0.03	0.01	0.00	0.00
16:1	11.61	12.22	10.08	5.57	12.50	7.76	10.41	10.66	2.84	3.51
17:1	1.37	1.27	0.47	0.28	0.31	0.30	1.68	1.65	0.07	0.08
18:1n9	9.22	11.94	7.62	8.59	10.27	10.04	10.85	10.14	10.65	8.70
20:1	0.20	0.04	0.00	0.05	0.00	0.04	0.10	0.02	0.10	0.07
22:1n9	0.03	0.00	0.00	0.06	0.00	0.00	0.00	0.00	0.01	0.01
24:1	0.16	0.22	0.00	0.00	0.00	0.00	0.00	0.40	0.04	0.00
MUFA	22.64	25.68	18.17	14.55	23.10	18.13	23.13	22.90	13.71	12.37
18:3 n 3	7.41	8.66	19.99	19.10	8.07	8.79	17.45	17.28	7.94	6.98
18:4 n 3	1.47	1.28	3.12	2.12	0.89	1.22	3.67	3.60	1.56	0.93
20:5 n 3	4.00	10.55	16.39	13.85	16.88	26.29	17.31	15.93	11.07	12.54
20:3 n 3	0.06	0.06	0.20	0.18	0.37	0.30	0.23	0.23	0.98	0.48
22:6 n 3	1.92	0.15	0.00	0.32	0.30	0.00	0.33	0.13	27.08	29.50
n3	14.86	20.70	39.70	35.57	26.51	36.60	39.00	37.18	48.63	50.41
18:2 n 6	5.03	6.66	4.73	7.06	4.34	3.03	3.20	4.46	4.44	3.62
18:3 n 6	0.15	0.72	0.38	0.22	0.19	0.17	0.46	0.57	0.07	0.07
20:3 n 6	0.03	0.00	0.00	0.07	0.02	0.00	0.11	0.58	0.31	0.20
20:4 n 6	1.48	1.92	0.81	1.23	3.89	2.88	2.53	4.49	2.54	2.75
22:2 n 6	2.42	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.13	0.00
n6	9.49	9.93	5.92	8.57	8.44	6.09	6.30	10.10	7.50	6.64
n3:n6	1.57	2.08	6.70	4.14	3.11	6.31	6.49	4.93	7.10	7.84
PUFA	25.02	31.09	45.62	44.42	35.25	42.94	45.30	47.30	56.22	57.11

608

609

610

611 Appendix 2. Annual biomass inputs of fatty acids from senesced vegetation.

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

Fatty acid	Annual Biomass Input (g/m ² /year)	
	Fresh Alder	Fresh hemlock
14:0	10.5	3.75
15:0	1.33	0.66
16:0	90.58	20.1
17:0	1.96	0.85
18:0	24.08	11.98
20:0	12.92	4.08
22:0	22.86	2.75
23:0	1.51	3.02
24:0	13.06	3.97
SAFA	178.82	51.17
14:1	1.23	0.32
15:1	0	0.81
16:1	5.18	2.38
17:1	0.46	0.84
18:1n9	44.38	14.76
20:1	1.05	0
22:1n9	0	0.26
24:1	0	0
MUFA	51.21	19.37
18:3 n 3	66.43	8.79
18:4 n 3	6.23	0.74
20:5 n 3	2.73	0
20:3 n 3	0.77	0.29
22:6 n 3	0.95	5.38
n3	77.11	15.20
18:2 n 6	38.15	11.20
18:3 n 6	0.32	0.32
20:3 n 6	0	0
20:4 n 6	1.54	0
22:2 n 6	0	0
n6	40.01	11.52
n3:n6	7.32	1.52
PUFA	119.70	28.92
SAFA: (MUFA+PUFA)	3.68	1.06