

# Managing macropods without poisoning ecosystems

By **Jordan O. Hampton** , **James M. Pay**, **Todd E. Katzner**, **Jon M. Arnemo**, **Mark A. Pokras**, **Eric Buenz**, **Niels Kanstrup**, **Vernon G. Thomas**, **Marcela Uhart**, **Sergio A. Lambertucci**, **Oliver Krone**, **Navinder J. Singh**, **Vinny Naidoo**, **Mayumi Ishizuka**, **Keisuke Saito**, **Björn Helander** and **Rhys E. Green**

*Jordan O. Hampton* is a Research Fellow in the Faculty of Veterinary and Agricultural Sciences at the University of Melbourne (Flemington Rd, Parkville, Vic. 3052, Australia; Tel: 0497 093 754; Email: jordan.hampton@unimelb.edu.au). *James M. Pay* is a post-doctoral Research Fellow in the School of Natural Sciences at the University of Tasmania (Churchill Ave, Hobart, Tas. 7005, Australia; Email: james.pay@utas.edu.au). *Todd E. Katzner* is a Research Wildlife Biologist in the Forest & Rangeland Ecosystem Science Center at the U.S. Geological Survey (230 N Collins Rd, Boise, ID 83702, USA; Email: tkatzner@usgs.gov). *Jon M. Arnemo* is a Professor of Biomedicine in the Department of Forestry and Wildlife Management at the Inland Norway University of Applied Sciences (NO-2480, Koppang, Norway) and Visiting Professor at the Swedish University of Agricultural Sciences (SE-90183 Umeå, Sweden; Email: jon.arnemo@inn.no). *Mark A. Pokras* is an Associate Professor Emeritus in the Cummings School of Veterinary Medicine, Tufts University (N. Grafton, MA, USA; Email: mark.pokras@tufts.edu). *Eric Buenz* is a Research Professor in the Nelson Marlborough Institute of Technology (NMIT) (322 Hardy Street, Nelson 7010, New Zealand; Email: eric.buenz@nmit.ac.nz). *Niels Kanstrup* is an Adjunct Associate Professor in the Department of Ecoscience at Aarhus University (Grenåvej 14, 8410, Ronde, Denmark; Email: nk@ecos.au.dk). *Vernon G. Thomas* is a Professor Emeritus in the Department of Integrative Biology at the University of Guelph (50 Stone Rd E, Guelph, ON N1G 2W1, Canada; Email: vthomas@uoguelph.ca). *Marcela Uhart* is Director, Latin America Program, in the One Health Institute, School of Veterinary Medicine, University of California (1089 Veterinary Medicine Drive, VM3B, Ground Floor, Davis, CA 95616, USA; Email: mubart@ucdavis.edu), *Sergio A. Lambertucci* is Principal Investigator in the Instituto de Investigaciones en Biodiversidad y Medio Ambiente (INIBIOMA) at National Scientific

**Summary** A recent review of the management of hyperabundant macropods in Australia proposed that expanded professional shooting is likely to lead to better biodiversity and animal welfare outcomes. While the tenets of this general argument are sound, it overlooks one important issue for biodiversity and animal health and welfare: reliance on toxic lead-based ammunition. Lead poisoning poses a major threat to Australia's wildlife scavengers. Current proposals to expand professional macropod shooting would see tonnes of an extremely toxic and persistent heavy metal continue to be introduced into Australian environments. This contrasts with trends in many other countries, where lead ammunition is, through legislation or voluntary programs, being phased out. Fortunately, there are alternatives to lead ammunition that could be investigated and adopted for improved macropod management. A transition to lead-free ammunition would allow the broad environmental and animal welfare goals desired from macropod management to be pursued without secondarily and unintentionally poisoning scavengers. Through this article, we hope to increase awareness of this issue and encourage discussion of this potential change.

**Key words:** animal welfare, culling, harvesting, One Health, scavengers, toxicology.

and Technical Research Council (CONICET) (Quintral 1250, R 8400 FRF, Bariloche, Argentina; Email: slambertucci@comabue-conicet.gob.ar). *Oliver Krone* is Head of Parasitology And Toxicology in the Department of Wildlife Diseases at the Leibniz Institute for Zoo and Wildlife Research (Alfred-Kowalke-Str. 17, 10315 Berlin, Germany; Email: krone@izw-berlin.de). *Navinder J. Singh* is Senior Lecturer & Associate Professor in the Department of Wildlife, Fish and Environmental Studies at the Swedish University of Agricultural Sciences (SE- 90183 Umeå, Sweden; Email: navinder.singh@slu.se). *Vinny Naidoo* is Dean in the Faculty of Veterinary Science at the University of Pretoria (Private Bag X04, Onderstepoort 0110, South Africa; Email: vinny.naidoo@up.ac.za). *Mayumi Ishizuka* is a Professor in the Department of Environmental Veterinary Sciences at Hokkaido University (9 Chome Kita 18 Jonishi, Kita Ward, Sapporo, Hokkaido 060-0818, Japan; Email: isbizum@vetmed.bokudai.ac.jp). *Keisuke Saito* is President of the Institute for Raptor Biomedicine Japan (2-2101, Hokuto, Kushiro-shi, Hokkaido 084-0922, Japan; E-mail: k\_saito@irbj.net). *Björn Helander* is a Senior scientist in Environmental Research & Monitoring at the Swedish Museum of Natural History

(Box 50007, SE-104 05 Stockholm, Sweden; Email: bjorn.belander@nrm.se). *Rhys E. Green* is Professor of Conservation Science in the Department of Zoology at the University of Cambridge (Pembroke Street, Cambridge CB2 3QZ, UK; Email: reg29@hermes.cam.ac.uk).

## Implications to Managers

- Lead is a toxic and persistent heavy metal that continues to be used in ammunition for macropod management in Australia.
- Lead fragments in the carcasses of shot macropods are ingested by scavenging wildlife, with negative effects on their survival, health and welfare.
- In recognition of the threat posed by lead ammunition to

wildlife and people, several nations have required or encouraged use of lead-free alternatives; this approach has not been discussed for macropods.

- Increased awareness of the threat posed by lead and the broad One Health benefits of lead-free alternatives is warranted in discussions of improvements to macropod management.

## Introduction

We applaud editors John L. Read, Graeme Coulson, James Q. Radford and George R. Wilson, and all contributing authors, for raising the profile of Australian macropod management, via their Special Issue of *Ecological Management & Restoration* 'Optimum management of overabundant macropods' (November, 2021). We agree with the main conclusions of this volume, that management of overabundant macropod populations is most effective if it is evidence-based, proactive and adaptive to minimise waste and use resources sustainably. We similarly appreciate that professional shooting of hyperabundant macropods, as opposed to alternative management options, is likely to lead to better biodiversity and animal welfare outcomes (Read *et al.* 2021). However, one important issue has been overlooked in this volume: all professional shooting of macropods currently relies on toxic lead-based ammunition.

Lead is a heavy metal that is toxic to nearly all life (Arnemo *et al.* 2016). In fact, anthropogenic lead pollution is a classical One Health problem: affecting the health of humans, animals and the environment (Hampton *et al.* 2018). Australians commonly eat meat from macropods killed with lead-based bullets, and there is a potential but so far unquantified risk to human health. This contrasts to elsewhere in the world where lead residues in game meat are a major concern and focus for research (Thomas *et al.* 2020). The problem we draw attention to here is the pronounced risk that

spent lead ammunition poses to scavengers. Scavengers play critical ecological roles in wildlife communities (Wilson & Wolkovich 2011) and are threatened by lead exposure worldwide (Pain *et al.* 2019). Tiny fragments are created by the frangible, 'varmint'-style, lead bullets typically used for macropod shooting (McTee *et al.* 2017). These fragments disperse widely through the tissues of shot animals (heads in the case of macropods; Figure 1), which are left in the field by shooters (Brooker & Ridpath 1980) and are readily ingested by scavengers (Figure 2) (except in the rare cases, where culled macropods are buried; Gordon 2019).


Avian, mammalian and reptilian scavengers feed extensively on discarded macropod tissues (Read & Wilson 2004). Acute lethal exposure can be widespread as a result of this process (Slabe *et al.* 2022) but equally concerning is the potential for sublethal effects resulting in chronic health degradation (Pay *et al.* 2021), as has been demonstrated for Golden Eagles (*Aquila chrysaetos*) in Europe (Ecke *et al.* 2017; Singh *et al.* 2021). Avian species known to be at-risk throughout Australia include Wedge-Tailed Eagles (*A. audax*) (Lohr *et al.* 2020; Hampton *et al.* 2021; Pay *et al.* 2021; Figure 2). In Tasmania, taxa documented as at-risk also include the threatened Tasmanian Devil (*Sarcophilus harrisi*) (Hivert *et al.* 2018). Furthermore, although data are sparse, a suite of other scavenging species in Australia (e.g. corvids; Figure 2) also likely encounter risk from lead poisoning when feeding on lead-killed carcasses.

The current scale of spent lead ammunition distributed on the landscape via macropod shooting is remarkable. More than 1.4 million macropods have been commercially harvested in Australia each year since 2010 (Department of Agriculture, Water and the Environment 2020). Through these actions, a conservative estimate is that 1.4 tonnes of lead fragments are introduced into the rangelands annually and consequently are available to scavengers (Hampton *et al.* 2018). Although spread over millions of hectares, this is still an immense quantity of lead, and this

estimate only considers commercial harvesting, not the extensive non-commercial culling that occurs for local population management. Furthermore, lead from ammunition also can persist for >30 years (Kanstrup *et al.* 2020), polluting soils, waterways and associated wildlife (Scheuhammer *et al.* 2003).

Although under-recognised in Australia, ecotoxicology is influential for wildlife populations (biodiversity) and individuals (animal health and welfare) in the Anthropocene (Death *et al.* 2019). Indeed, the issue of lead poisoning resulting from spent lead ammunition is rarely mentioned in the ecological or conservation communities within Australia (Hampton *et al.* 2018). The negative impacts of lead are also not mentioned in recent reviews of animal welfare and kangaroo management (McLeod & Sharp 2014; Descovich *et al.* 2015; Wilson & Edwards 2019; Stephens 2021). In addition to direct impacts (Stephens 2021), a complete assessment of animal welfare impacts would be expected to include consideration of non-target animals affected by indirect processes such as secondary lead poisoning (Hampton *et al.* 2019), as increasingly occurs for other wildlife management issues, such as use of anticoagulant rodenticides (Fisher *et al.* 2019).

All of this is in stark contrast to what is happening outside Australia. In recognition of the threat posed to wildlife and people, programs to reduce use of lead ammunition are expanding globally. While lead ammunition (shotgun shot) has been banned for waterfowl hunting in many Australian jurisdictions for decades, legislation has not addressed the widespread use of shotgun in other contexts or any use of rifle bullets (Hampton *et al.* 2018). In contrast, the US state of California recently banned use of lead rifle bullets (Schulz *et al.* 2021). More recently, Denmark has announced a ban of all lead ammunition for hunting by 2023 (Kanstrup *et al.* 2021), and the European Union (Thomas *et al.* 2021), United Kingdom (UK Government 2021) and Japan (Ishii *et al.* 2020) all are considering or have passed regulations to do the same. Likewise, voluntary programs to encourage use of lead-free ammunition are



**Figure 1.** Radiographs (X-rays) of the heads of Western Grey Kangaroos (*Macropus fuliginosus*) shot with lead-based bullets by a commercial harvester. Bright white lead fragments (some shown by red arrows) are seen distributed widely through each head. These tiny and toxic fragments are easily ingested by wildlife scavengers such as raptors.



**Figure 2.** Avian scavengers of shot macropod carcasses include (a) Wedge-Tailed Eagles (*Aquila audax fleayi*) with a Tasmanian Pademelon (*Thylogale billardieri*) and (b) Australian Ravens (*Corvus* spp.) feeding on a Western Grey Kangaroo shot in mainland Australia.

becoming more abundant (Schulz *et al.* 2021). Finally, these actions are also affecting private industry overseas – for example, the major U.K. game-retailing supermarket no longer stocks meat from animals shot with lead ammunition (Thomas *et al.* 2020).

The absence of such legislation or other action to reduce lead exposure of scavengers in Australia is certainly not the fault of macropod managers. However, the management plans proposed in the recent Special Issue ‘Optimum management of overabundant macropods’ would be improved by consideration of animals affected by indirect processes such as heavy metal pollution (Hampton *et al.* 2019). In fact, it seems that any proposal that fails to consider the profound impacts of lead pollution is unlikely to represent an ‘optimum’ approach. Likewise, procedural guidelines for shooting macropods would be improved by considering this process. There is, surprisingly, no discussion of lead or the role of lead-

free ammunition in the recently updated ‘National Code of Practice for the Humane Shooting of Kangaroos and Wallabies for Commercial Purposes’ (Agrifutures Australia 2020).

Use of lead ammunition pollutes the environment and poisons Australia’s scavengers. However, there are alternatives that allow for exactly the same management actions without the secondary consequences to local wildlife or public health. The proposals of Read *et al.* (2021), as currently stated, would perpetuate the pollution of environments with high conservation value with tonnes of an extremely toxic and persistent heavy metal. It is apparent that there is currently limited consideration of this issue in Australian wildlife management. An important goal of this article is to increase awareness and consideration of this issue.

There is an ever-increasing range of alternatives to lead-based bullets being used globally, mostly being copper-based (Kanstrup & Thomas 2020). However,

there are some special challenges associated with finding appropriate lead-free bullets for macropod shooting, aside from generally higher cost (Hampton *et al.* 2020). Relatively small calibres ( $\leq 0.223$ ) are typically used, with important interactions between bullet length and barrel twist rates that affect precision (Caudell *et al.* 2012), especially when considering the minimum bullet weights specified by the commercial macropod shooting Code of Practice (Agrifutures Australia 2020). Nonetheless, rimfire and centrefire (0.223) lead-free bullets have been shown to produce comparable outcomes for the shooting of other mammalian species (McTee *et al.* 2017). Promoting a transition to lead-free ammunition for macropod management may be more challenging than for contexts such as deer hunting, where abundant information can be drawn from international studies (Stokke *et al.* 2019). There is, therefore, value to structured studies assessing the different types of lead-free bullets for



specific macropod shooting requirements to ensure that efficacy and animal welfare standards can be maintained (Hampton *et al.* 2020). Once that step is taken, shooters will have the option to transition to non-toxic products and some of the broad environmental and animal welfare goals desired from macropod management may be pursued without secondarily and unintentionally poisoning scavengers.

## Acknowledgements

This article was written in response to the special issue 'Optimum management of overabundant macropods' published in *Ecological Management & Restoration* (November, 2021). The writing of this manuscript received no specific funding. We acknowledge the contributions of D. Pain and D. Forsyth to the initial stages of this manuscript. Any use of trade, firm or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. Open access publishing facilitated by The University of Melbourne, as part of the Wiley - The University of Melbourne agreement via the Council of Australian University Librarians.

## References

- Agrifutures Australia (2020) *National Code of Practice for the Humane Shooting of Kangaroos and Wallabies for Commercial Purposes*. AgriFutures Australia, Wagga Wagga.
- Arnemo J. M., Andersen O., Stokke S. *et al.* (2016) Health and environmental risks from lead-based ammunition: science versus socio-politics. *EcoHealth* **13**, 618–622.
- Brooker M. G. and Ridpath M. (1980) The diet of the wedge-tailed eagle, *Aquila audax*, in Western Australia. *Wildlife Research* **7**, 433–452.
- Caudell J. N., Stopak S. R. and Wolf P. C. (2012) Lead-free, high-powered rifle bullets and their applicability in wildlife management. *Human-Wildlife Interactions* **6**, 105–111.
- Death C. E., Griffiths S. R. and Story P. G. (2019) Terrestrial vertebrate toxicology in Australia: an overview of wildlife research. *Current Opinion in Environmental Science and Health* **11**, 43–52.
- Department of Agriculture, Water and the Environment (2020) *Macropod Quotas and Harvest Statistics for NSW, QLD, SA and WA – 2020*. Department of Agriculture, Water and the Environment, Canberra.
- Descovich K., McDonald I., Tribe A. and Phillips C. (2015) A welfare assessment of methods used for harvesting, hunting and population control of kangaroos and wallabies. *Animal Welfare* **24**, 255–265.
- Ecke F., Singh N. J., Arnemo J. M. *et al.* (2017) Sublethal lead exposure alters movement behavior in free-ranging golden eagles. *Environmental Science & Technology* **51**, 5729–5736.
- Fisher P., Campbell K. J., Howald G. R. and Warburton B. (2019) Anticoagulant rodenticides, islands, and animal welfare accountability. *Animals* **9**, 919.
- Gordon I. J. (2019) Adopting a utilitarian approach to culling wild animals for conservation in National Parks. *Conservation Science and Practice* **1**, e105.
- Hampton J. O., DeNicola A. J. and Forsyth D. M. (2020) Assessment of lead-free, 22 LR bullets for shooting European rabbits. *Wildlife Society Bulletin* **44**, 760–765.
- Hampton J. O., Laidlaw M., Buenz E. and Arnemo J. M. (2018) Heads in the sand: public health and ecological risks of lead-based bullets for wildlife shooting in Australia. *Wildlife Research* **45**, 287–306.
- Hampton J. O., Specht A. J., Pay J. M., Pokras M. and Bengsen A. J. (2021) Portable X-ray fluorescence for bone lead measurements of Australian eagles. *Science of the Total Environment* **789**, 147998.
- Hampton J. O., Warburton B. and Sandøe P. (2019) Compassionate versus consequentialist conservation. *Conservation Biology* **33**, 751–759.
- Hivert L., Clarke J., Peck S. *et al.* (2018) High blood lead concentrations in captive Tasmanian devils (*Sarcophilus harrisii*): a threat to the conservation of the species? *Australian Veterinary Journal* **96**, 442–449.
- Ishii C., Ikenaka Y., Nakayama S. M. *et al.* (2020) Current situation regarding lead exposure in birds in Japan (2015–2018); lead exposure is still occurring. *Journal of Veterinary Medical Science* **82**, 1118–1123.
- Kanstrup N., Balsby T. J. S., Møllerup K. A. and Hansen H. P. (2021) Non-lead rifle ammunition: Danish hunters' attitudes. *Environmental Sciences Europe* **33**, 41.
- Kanstrup N., Fox A. D. and Balsby T. J. S. (2020) Toxic lead gunshot persists accessible to waterbirds after a 33-year ban on their use. *Science of the Total Environment* **714**, 136876.
- Kanstrup N. and Thomas V. G. (2020) Transitioning to lead-free ammunition use in hunting: socio-economic and regulatory considerations for the European Union and other jurisdictions. *Environmental Sciences Europe* **32**, 91.
- Lohr M. T., Hampton J. O., Cherriman S., Busetti F. and Lohr C. (2020) Completing a worldwide picture: preliminary evidence of lead exposure in a scavenging bird from mainland Australia. *Science of the Total Environment* **715**, 135913.
- McLeod S. R. and Sharp T. M. (2014) *Improving the Humaneness of Commercial Kangaroo Harvesting*. Rural Industries Research and Development Corporation, Canberra.
- McTee M., Young M., Umansky A. and Ramsey P. (2017) Better bullets to shoot small mammals without poisoning scavengers. *Wildlife Society Bulletin* **41**, 736–742.
- Pain D. J., Mateo R. and Green R. E. (2019) Effects of lead from ammunition on birds and other wildlife: a review and update. *Ambio* **48**, 935–953.
- Pay J. M., Katzner T. E., Hawkins C. E. *et al.* (2021) High frequency of lead exposure in the population of an endangered Australian top predator, the Tasmanian wedge-tailed eagle (*Aquila audax fleayi*). *Environmental Toxicology and Chemistry* **40**, 219–230.
- Read J. and Wilson D. (2004) Scavengers and detritivores of kangaroo harvest offcuts in arid Australia. *Wildlife Research* **31**, 51–56.
- Read J. L., Wilson G. R., Coulson G. *et al.* (2021) Improving kangaroo management: a joint statement. *Ecological Management & Restoration* **22**, 186–192.
- Scheuhammer A. M., Bond D. E., Burgess N. M. and Rodrigue J. (2003) Lead and stable lead isotope ratios in soil, earthworms, and bones of American woodcock (*Scolopax minor*) from eastern Canada. *Environmental Toxicology and Chemistry* **22**, 2585–2591.
- Schulz J. H., Stanis S. A. W., Hall D. M. and Webb E. B. (2021) Until it's a regulation it's not my fight: complexities of a voluntary nonlead hunting ammunition program. *Journal of Environmental Management* **277**, 111438.
- Singh N. J., Ecke F., Katzner T., Bagchi S., Sandström P. and Hörnfeldt B. (2021) Consequences of migratory coupling of predators and prey when mediated by human actions. *Diversity and Distributions* **27**, 1848–1860.
- Slabe V. A., Anderson J. T., Millsap B. A. *et al.* (2022) Demographic implications of lead poisoning for eagles across North America. *Science* **375**, 779–782.
- Stephens T. (2021) Kangaroo management and animal welfare. *Ecological Management and Restoration* **22**, 71–74.
- Stokke S., Arnemo J. M. and Brainerd S. (2019) Unleaded hunting: are copper bullets and lead-based bullets equally effective for killing big game? *Ambio* **48**, 1044–1055.
- Thomas V. G., Kanstrup N. and Pain D. J. (2021) Promoting the transition to non-lead hunting ammunition in the European Union through regulation and policy options. *Environmental Policy and Law* **51**, 239–254.
- Thomas V. G., Pain D. J., Kanstrup N. and Green R. E. (2020) Setting maximum levels for lead in game meat in EC regulations: an adjunct to replacement of lead ammunition. *Ambio* **48**, 954–968.
- UK Government (2021) *Plans Announced to Phase out Lead Ammunition in Bid to Protect Wildlife*. UK Government, London.
- Wilson E. E. and Wolkovich E. M. (2011) Scavenging: how carnivores and carrion structure communities. *Trends in Ecology and Evolution* **26**, 129–135.
- Wilson G. R. and Edwards M. (2019) Professional kangaroo population control leads to better animal welfare, conservation outcomes and avoids waste. *Australian Zoologist* **40**, 181–202.