

Peripheral adaptations to endurance training—Effect of active muscle mass

Knut Sindre Mølmen^{1,2}  | Jostein Hallén¹ | Bjarne Rud¹

¹Department of Physical Performance, Norwegian School of Sport Sciences, Oslo, Norway

²Section for Health and Exercise Physiology, Inland Norway University of Applied Sciences, Lillehammer, Norway

Correspondence

Knut Sindre Mølmen, Inland Norway University of Applied Sciences, PO. Box 422, 2604 Lillehammer, Norway.
Email: knut.sindre.molmen@inn.no

Abstract

The purpose of this study was to evaluate whether adaptations to endurance training are affected by active muscle mass during training. Eleven healthy subjects performed 5 weeks of one-legged knee extension (1-KE) training with both legs, separately. During the 1-KE workouts for one of the legs, arm cycling was added (1L2A), while the other leg only performed 1-KE (1L). During the training sessions, the 1-KE power output and session duration were equal for both legs. Whole-body oxygen uptake (VO_2), adrenaline, and noradrenaline plasma concentrations were $112\% \pm 11\%$, $139\% \pm 144\%$, and $197\% \pm 101\%$ higher during 1L2A than 1L, respectively. However, this did not affect the 1-KE training adaptations since submaximal O_2 -cost, heart rate and blood lactate concentration, maximal VO_2 , and power output all improved equally in both legs. This was supported by similar capillarization and concentration of oxidative enzymes in the legs after the training period. The present study therefore indicates that the size of active muscle mass per se during exercise does not affect adaptations to endurance training.

KEYWORDS

arm cranking, muscle fiber, pulmonary oxygen uptake, single-leg, skeletal muscle, training adaptation

1 | INTRODUCTION

Whole-body endurance training potentially leads to both central and peripheral adaptations. The peripheral adaptations include muscular capillarization and increased oxidative capacity.¹ However, how the peripheral adaptations are regulated is not fully understood and there are contradicting findings regarding the importance of different factors such as high mass-specific skeletal muscular energy turnover and high local metabolic stress. For example, comparable endurance training studies (ie, similar subject and training characteristics) with one-legged exercise and whole-body exercises exhibit approximately the same skeletal muscular adaptations. Typically 20%-30% increased capillarization and 20%-40% increases in oxidative enzymes activity after

6-8 weeks of training are reported.²⁻⁵ This happens though one-legged exercises clearly provide a higher available capacity for mass-specific skeletal muscular energy turnover than whole-body exercises.^{6,7} On the other hand, when exercising with a large muscle mass, the sympathetic response is higher than when exercising with a small muscle mass.⁸ This is potentially beneficial for muscular adaptations.

It has previously been shown that treatment with β -adrenergic blockers attenuates the acute increase in gene expression of the transcriptional coactivator peroxisome proliferator-activated receptor- γ coactivator (PGC-1 α), known as a key regulator of mitochondrial biogenesis, after endurance exercise.^{9,10} In addition, β -adrenergic blockers seem to hamper the mitochondrial adaptations to endurance training.^{11,12} However, these studies are also accompanied by contradictory findings.¹³⁻¹⁵

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors *Translational Sports Medicine* Published by John Wiley & Sons Ltd.

Brandt et al¹⁶ compared acute responses to matched cycling workloads (corresponding to 60% VO₂max) for 60 minutes with or without the addition of 1-minute bouts of intermittent arm cycling every 5 minutes (~100 watts of additional workload). The addition of arm cycling resulted in a higher plasma adrenaline concentration. However, the PGC-1 α mRNA response was similar. The same was the phosphorylation of upstream targets assumed to upregulate PGC-1 α (ie, cAMP-response element binding, p38 mitogen-activated protein kinase [p38 MAPK], and AMP-activated protein kinase [AMPK]). Somewhat contrary to this finding did Fiorenza et al¹⁷ observe a higher exercise-induced phosphorylation of p38 MAPK and Ca²⁺/calmodulin-dependent protein kinase II (CAMKII) after speed endurance exercise compared with repeated sprints and high-volume continuous moderate-intensity cycling. The speed endurance protocol was associated with greater catecholamine response than the other two protocols, in addition to the most marked metabolic perturbations as evidenced by the greatest changes in muscle lactate and pH. However, the PGC-1 α mRNA response was similar between speed endurance and continuous cycling, while it was lower after repeated sprint exercise.

Whether differences in non-pharmacologically induced sympathetic response per se can influence skeletal muscular adaptations to endurance training in humans has not previously been reported. Abbiss et al¹⁸ found higher improvement in markers of oxidative enzymes (ie, cytochrome *c* oxidase subunit II and IV protein concentration) after 3 weeks of one-legged cycling compared to bicycling, indicating that training with a small active muscle mass is more favorable for local muscular adaptations compared to whole-body exercise. However, because leg workload, and thus leg energy turnover, was higher during one-legged cycling (13%), such a study design is not appropriate for evaluating the importance of active muscle mass per se since not all confounding factors are taken into account. The study of MacInnis et al¹⁹ compared training adaptations of work-matched one-legged moderate-intensity continuous and high-intensity interval cycling training. They concluded that the high-intensity interval protocol elicited superior mitochondrial adaptations, due to the greater increase in citrate synthase activity and mass-specific oxidative phosphorylation capacities than continuous exercise. The high-intensity exercise probably induced a higher catecholamine response due to the higher exercise intensity, but it probably also induced greater metabolic perturbations as seen in Fiorenza et al,¹⁷ which makes it difficult to evaluate causality in the training adaptations.

The purpose of the present study was therefore to test whether local training adaptations are affected by the size of the active muscle mass per se during exercise. This was evaluated by adding arm cycling to one-legged knee extension (1-KE) for one of the legs only, while the other leg was kept as a control, performing ordinary 1-KE. Previously, it has

been shown that O₂ delivery to the exercising leg in healthy subjects is maintained during combined arm cycling and 1-KE.^{8,20} Therefore, with this model the legs can exercise with equal power output and energy demand. The maintained oxygen supply in this model should ensure similar metabolic stress in the exercising leg muscles even with the addition of arm cycling. However, whole-body energy turnover and catecholamine response will most likely increase with added arm cycling. We hypothesized that the addition of arm cycling to the 1-KE exercise would elicit greater training adaptations.

2 | METHODS

Eleven healthy subjects (two females) provided informed consent to participate in the study (Table 1). The project was approved by Norwegian Regional Committee for Medical Research Ethics and performed according to the Declaration of Helsinki. The subjects were familiarized to the testing and training protocol over a minimum of 4 days to exclude familiarization effects. They were instructed to maintain their usual physical activity level during the study and abstain from strenuous exercise 24 hours prior to the experimental trials.

2.1 | Training protocol

Each subject took part in 3- to 4-weekly training sessions for 5 weeks. The training consisted of dynamic 1-KE with a frequency of 60 contractions per min, performed separately on the subject's left and right legs using a custom-designed electromagnetic braked ergometer²¹ that addressed external work to only the *quadriceps femoris* muscle.²² One of the legs (randomized for left or right leg) was trained simultaneously with two-arm cycling (1L2A) (Excalibur Sport; Lode BV), while the arms were kept relaxed when the control leg was trained (1L; Figure 1). However, the workloads of the legs were equal. The two training exercises were performed in the same session with a 10-minute break between sessions to avoid any carry-over sympathetic response effect, as catecholamines have a half-life in plasma of ~2 minutes.²³ The training sessions therefore consisted of two 20- to 30-minute bouts (gradually increased from 20-minute bouts in week

TABLE 1 Subjects' characteristics

Age (years)	27 ± 5
Height (cm)	179 ± 8
Mass (kg)	80 ± 11
Body mass index	25 ± 2
VO ₂ max (mL kg ⁻¹ min ⁻¹)	52 ± 6

Note: Values presented as mean ± standard deviation. VO₂max, maximal oxygen uptake (pulmonary) (n = 11).




FIGURE 1 The training exercises 1L (left) and 1L2A (right)

1, 25-minutes bouts in weeks 2-3, and 30-minutes bouts in weeks 4-5), one being 1L and the other being 1L2A. The workloads were individually adjusted daily, with the aim of nearly exhausting the subjects at the end of the exercise bout, but always with the same power and duration for both legs during the same workout. This corresponded to an average 1-KE exercise intensity of $73\% \pm 3\%$ of baseline peak power output (range, 67%-78%), while the arm cycling exercise intensity during 1L2A was $54\% \pm 5\%$ (range, 45%-61%) of arm cranking peak power output. Start-up exercise was counter-balanced between 1L and 1L2A in the way that left and right legs were exercised first every second time.

During two separate workouts in the latter part of the training period, blood samples were taken immediately after finishing the first exercise bout of the workout to examine venous blood plasma concentrations of adrenaline and nor-adrenaline. The subjects arrived at the laboratory in the morning after an overnight fast or in the afternoon having fasted for 4 hours before training. Vacutainer[®] glass tubes (Becton Dickinson) with added EGTA-glutathione were used for the blood collection. The glass tubes were pre-treated at the Section of Specialized Endocrinology (Clinic of Medicine, Oslo University Hospital) and later analyzed there. During exercise, measurements of pulmonary oxygen consumption (VO_2), ventilation, and respiratory exchange ratio were carried out using an automated system (JAEGER Oxycon PRO[™]; Carefusion GmbH), which was calibrated according to the instruction manual. In addition, heart rate was monitored continuously with a Polar monitor (Polar Electro Oy) and a fingertip capillary blood sample was analyzed for lactate concentration after completing the exercise (YSI 1500 Sport; Yellow Springs Instruments).

2.2 | Exercise tests

Prior to and after the training period, a set of exercise tests including bicycling, 1-KE, and arm cycling were performed.

Maximal pulmonary oxygen uptake ($\text{VO}_{2\text{max}}$) was tested on a bicycle ergometer (Excalibur Sport; Lode BV). A stepwise incremental protocol was used, and the initial power output

was individually set to elicit exhaustion after approximately 6 minutes. Power output was increased by 20 or 25 watts (W) each minutes for women and men, respectively. Exhaustion was defined as when the subject could not maintain a cadence of 80 revolutions per min (RPM). VO_2 , ventilation, respiratory exchange ratio, and heart rate were measured continuously, while tests of capillary lactate concentration and ratings of perceived exertion²⁴ were completed afterward. Time to exhaustion was noted during all maximal tests (mm:ss).

On a separate day, the subjects underwent 1-KE and arm cycling tests. Initially, three 5-minutes submaximal 1-KE workloads were completed (at 20, 30, and 40 W). Thereafter, a maximal 1-KE stepwise incremental protocol of 5 W increase per min, starting at 20 W, was conducted on the same leg until exhaustion, which was defined as the inability to maintain the contraction frequency (60 RPM). After resting for 90 minutes, the tests were repeated with the subjects' contralateral legs. Finally, they did a stepwise incremental arm cycling protocol, starting at 30 or 40 W, and increased by 10 or 15 W every minute until exhaustion (RPM < 60), for women and men, respectively. VO_2 , ventilation, respiratory exchange ratio, and heart rate were measured continuously during all the tests, while measurements of capillary lactate concentration and ratings of perceived exertion were completed immediately after exhaustion. During maximal 1-KE, some subjects exhibited a more elevated VO_2 at exhaustion than expected from the increases in workload, despite the subjects being fastened tightly with a four-point seat belt and trained to keep their non-exercising extremities as relaxed as possible. In accordance with Andersen et al,²² we therefore corrected $\text{VO}_{2\text{max}}$ by extrapolating the VO_2 /power output relationship from submaximal values for all subjects. All the tests were conducted in the same environment with the same protocol, equipment, and test personnel.

2.3 | Biopsy

After the training period, muscle biopsies of the mid-portion of the *vastus lateralis* muscle were taken from the subjects'

1L- and 1L2A-trained legs under local anesthesia, as described in detail elsewhere.²⁵

2.3.1 | Histochemical analysis

Initially, 8- μ m cross sections from the biopsies were incubated for 30 minutes in BSA blocking solution (10% BSA in PBS, diluted 1:6 in deionized water; Thermo Fisher Scientific, Inc). Subsequently, they were incubated with primary antibodies against muscle fiber membrane (rabbit anti-dystrophin, diluted 1:500; Abcam plc[®]), myosin heavy chain type II (mouse anti-SC71, diluted 1:100; DSHB), and capillaries (mouse anti-CD31, diluted 1:100; Dako). Anti-dystrophin and anti-SC71 were incubated at room temperature for 60 minutes, while anti-CD31 was incubated overnight at 4°C, followed by incubation with appropriate secondary antibodies (Alexa 488 or 494; Biotium Inc) for 60 minutes at room temperature. Between stages, sections were washed for 3 \times 10 minutes in 0.05% PBS-t solution. The stained sections were afterward visualized using a high-resolution camera (DP72; Olympus Corp.) mounted on a microscope (BX61; Olympus Corp.) with a fluorescent light source (X-Cite 120PCQ; EXFO Photonic Solutions Inc). Fiber type distribution, fiber cross-sectional area, and capillaries were identified using TEMA software (CheckVision). Capillarization was expressed as capillaries around each fiber (CAF), and CAF was related to fiber area (CAFA) for type I and type II (IIa and IIx gathered) fibers, in addition to capillary density (capillaries per mm²) and capillaries per fiber (C/F).

2.3.2 | Protein analysis

Muscle samples were homogenized in 1 mL tissue protein extraction reagent (T-PER[®], Thermo Fisher Scientific, Inc) and 10 μ L Calbiochem[®] Protease Inhibitor Cocktail Set IV (Merck Millipore) per ~50 mg muscle tissue. The lysate was centrifuged at 10 000 g for 10 minutes at 4°C. Protein concentration was assessed with a commercial kit (Bio-Rad RC/DC[™] Protein Assay kit; Bio-Rad Laboratories, Inc), a filter photometer (Expert 96; ASYS Hitech GmbH) and the software provided (Kim version 5.45.0.1; Daniel Kittrich).

2.3.3 | Western blotting

Equal amounts of protein (35 μ g) per sample were separated on Mini-PROTEAN[®] TGX Stain-Free[™] gels (Bio-Rad Laboratories, Inc) at 200 volts for ~30 minutes and subsequently transferred to polyvinylidene difluoride membranes (Bio-Rad Laboratories, Inc) through blotting. Membranes were blocked with 5% nonfat milk solution (in Tris-buffered saline with 0.1% Tween-20; TBS-t) for 120 minutes. Afterward, primary antibodies for citrate synthase (rabbit anti-CS, diluted 1:2 000; Abcam plc[®]), cytochrome c oxidase

subunit 4 (mouse anti-COX4, diluted 1:1 000; Abcam plc[®]), and hydroxyacyl-coenzyme A dehydrogenase (rabbit anti-HADH; diluted 1:8 000; Abcam plc[®]) were incubated overnight at 4°C. Membranes were then washed for 15 minutes with TBS-t and 3 \times 5 minutes with Tris-buffered saline, followed by incubation with secondary antibodies to anti-mouse (goat anti-mouse, diluted 1:30 000; Thermo Fisher Scientific, Inc) or anti-rabbit (goat anti-rabbit, diluted 1:3 000; Cell Signaling) for 60 minutes at room temperature. All antibodies were diluted in a 1% fat-free skimmed milk and 0.05% TBS-t solution. After a new washing round and incubation with SuperSignal[®] West Dura Extended Duration Substrate (Thermo Fisher Scientific, Inc), the proteins were visualized by chemiluminescence and quantified by densitometry (ChemiDoc[™] MP; Bio-Rad Laboratories, Inc). All densitometry values were expressed relative to the total amount of protein (Bio-Rad stain-free method). All samples were run as duplicates, and mean values were used for statistical analyses. The 1L-trained leg was set as the control and 100%.

2.4 | Statistical analysis

Power calculations were performed to estimate the sample size needed to obtain an average increase of 0.2 L min⁻¹ in 1KE $\dot{V}O_{2\max}$ in 1L and an additional 0.075 \pm 0.075 L min⁻¹ in 1L2A. A sample size of 11 for each training modality should be sufficient to detect differences between legs (α level of 0.05). Variables with pre- and post-measurements for 1L- and 1L2A-trained legs were tested for mean differences with two-way repeated measures ANOVA with time (pre and post) and leg (1L and 1L2A) as within factors. When significant main effects were indicated, Fisher's least significant difference *post-hoc* tests were used to compare means. Differences and changes for the other variables studied were evaluated using paired two-tailed Student's *t* test (level of significance, $P < 0.05$). All results are presented as mean \pm standard deviation.

3 | RESULTS

Before training, the two legs were similar concerning all variables measured at that time point (Table 3 and Figure 2).

3.1 | Training

The participants performed 17.3 \pm 1.7 training sessions with 1L and 1L2A. Power output and amount of work were the same in both legs (in average, 35 \pm 8 W and 50 \pm 17 kJ per training session, respectively). Acute systemic responses, as evaluated by cardiorespiratory variables, capillary blood lactate concentrations, and venous blood catecholamine concentrations, were higher during 1L2A workouts than in 1L workouts (Table 2).




FIGURE 2 Submaximal 1-KE pre- and post-training values for O_2 -cost (A and B), HR (heart rate; C and D), and $[BLa^-]$ (capillary lactate concentration; E and F) for 1L and 1L2A. *, Significant different from pre-value ($P < 0.05$)

3.2 | Training effects

1-KE $\dot{V}O_{2\max}$ increased with training only for 1L2A ($P < 0.05$; Table 3). However, there was no interaction effect between the legs (1L2A vs 1L, $P = 0.52$). The same was true for 1KE $\dot{V}E_{\max}$ (interaction effect between 1L2A and 1L, $P = 0.33$). Peak power output was improved for both legs ($P < 0.05$), but in a similar manner ($P = 0.61$).

Capillary lactate concentration was reduced with training ($-21\% \pm 16\%$; $P < 0.05$), but no interaction effect was found between the two legs ($P = 0.18$, Figure 2). Submaximal 1-KE O_2 -cost ($-6\% \pm 10\%$; $P = 0.07$) and heart rate ($-6\% \pm 9\%$; $P = 0.05$) only tended to decrease with training.

$\dot{V}O_{2\max}$ during ordinary cycling and arm cycling improved during the training period by $3 \pm 3\%$ (from 52.4 ± 6.3 to 54.2 ± 6.7 $\text{mL kg}^{-1} \text{min}^{-1}$) and $8\% \pm 8\%$ (from 2310 ± 632 to 2482 ± 631 mL min^{-1}), respectively ($P < 0.05$ for both).

3.3 | Muscle fiber characteristics (after training)

Differences in muscle characteristics between the two legs were evaluated from biopsies taken after the training period. There were no differences in relative proportions

of fiber types, cross-sectional area, capillarization, or oxidative enzymes between the two legs (Table 4 and Figure 3).

TABLE 2 Acute physiological responses to the training workouts

	1L	1L2A
Power output, leg (W)	36 ± 9	36 ± 9
Power output, arms (W)	–	74 ± 18
NA (pmol L^{-1})	4465 ± 1866	12445 ± 3905^a
A (pmol L^{-1})	1341 ± 837	2462 ± 1161^a
$[BLa^-]$	2.7 ± 0.8	4.7 ± 1.2^a
HR (beats min^{-1})	108 ± 21	157 ± 15^a
$\dot{V}O_2$ (L min^{-1})	1.1 ± 0.3	2.3 ± 0.5^a
$\dot{V}E$ (L min^{-1})	33 ± 8	74 ± 14^a
RER	0.93 ± 0.06	0.96 ± 0.04
RPE	14.4 ± 1.6	14.4 ± 1.2

Note: Variables tested during an exercise session in the latter part of the training period (training session #13 out of 17; $n = 9$). Values are means \pm standard deviation. Noradrenaline (NA) and adrenaline (A) concentrations in venous blood plasma.

Abbreviations: $[BLa^-]$, capillary lactate concentration; HR, heart rate; RER, respiratory exchange ratio; RPE, rating of perceived exertion; $\dot{V}E$, ventilation; $\dot{V}O_2$, oxygen consumption (pulmonary).

^a P -value < 0.05 between exercises.

TABLE 3 Values achieved during maximal 1-KE

Exercise	VO ₂ max (L min ⁻¹)	VE _{max} (L min ⁻¹)	HR _{max} (beats min ⁻¹)	[BLa ⁻] (mmol L ⁻¹)	Peak power (W)
1-KE max					
1L pre	1.25 ± 0.26	62 ± 18	150 ± 21	3.8 ± 1.1	49 ± 12
1L post	1.31 ± 0.33	69 ± 22	151 ± 22	3.9 ± 0.8	55 ± 12 ^a
Change (%)	4 ± 9	11 ± 17	1 ± 10	8 ± 28	14 ± 7
1L2A pre	1.20 ± 0.24	57 ± 15	149 ± 19	3.7 ± 0.9	48 ± 12
1L2A post	1.29 ± 0.24 ^a	69 ± 24 ^a	147 ± 16	3.9 ± 0.8	55 ± 13 ^a
Change (%)	8 ± 10	21 ± 22	-1 ± 9	11 ± 26	16 ± 9

Note: Values are means ± standard deviation.

Abbreviations: [BLa⁻], capillary lactate concentration; HR, heart rate; VE, ventilation; VO₂, oxygen uptake (pulmonary); W, watt.

*P-Value <0.05 between pre- and post-values (n = 11).

4 | DISCUSSION

The present study shows that 5 weeks of 1-KE training with or without added arm training led to the same training adaptations demonstrated by similar improvements in 1-KE VO₂max and peak power output. In addition did the O₂-cost, heart rate, and capillary lactate concentration during submaximal 1-KE change in a similar manner, and capillarization and concentration of oxidative enzymes in the legs were matching after the training period. This occurred though whole-body aerobic energy turnover, blood lactate concentration, and catecholamine responses were different during the training workouts, which implies that the difference in systemic variables per se did not affect the training adaptations to endurance training.

There are not many studies comparing adaptations after training periods with exercises engaging different active muscle mass. To the field of exploring which factors that are important for training adaptations, the present study

indicates that systemic variables such as catecholamine response, capillary lactate levels, and whole-body aerobic energy turnover per se during exercise are probably not mandatory for muscular adaptations. It is therefore reasonable to assume that local evoked factors are of greater importance. Theoretically, these local factors could include the mass-specific energy turnover, either per se or in concert with metabolic stress. Metabolic stress with intramuscular metabolite accumulation of, that is, lactate, inorganic phosphate, and hydrogen ions, is associated with activation of metabolic pathways such as AMPK, CAMKII, p38 MAPK, and PGC-1 α that is important for muscular training adaptations.²⁶ This approach fits well with the previous findings of similar muscular adaptations after training with a small active muscle mass compared to whole-body exercises²⁻⁵ and with the finding of superior muscular adaptations in training protocols developing the highest metabolic stress, although the protocols were work-matched.²³ Blood flow occlusion during 1-KE training, and therefore restricted O₂ availability, has been shown to enhance the

TABLE 4 Muscle fiber characteristics

	1L	1L2A
Fiber type, fiber area, and capillarization		
Fiber type 1 (%)	45 ± 11	40 ± 8
CSA (μm^2) fiber type 1	5075 ± 1209	4251 ± 1238
CAF fiber type 1	5.6 ± 1.1	5.2 ± 1.1
CAFA fiber type 1	1.19 ± 0.23	1.31 ± 0.19
CSA (μm^2) fiber type 2	5347 ± 1433	4936 ± 1642
CAF fiber type 2	5.2 ± 0.8	5.1 ± 1.0
CAFA fiber type 2	1.05 ± 0.27	1.10 ± 0.22
CD (capillaries·mm ²)	494 ± 120	540 ± 100
C/F	2.49 ± 0.47	2.43 ± 0.53

Note: Values are means ± standard deviation.

Abbreviations: C/F, capillaries per fiber (n = 8); CAF, average number of capillaries around fiber; CAFA, CAF related to CSA; CD, capillary density; CSA, cross-sectional area.




FIGURE 3 Relative concentrations of oxidative enzymes. 1L leg set to 100%. CS, citrate synthase; HADH, hydroxyacyl-coenzyme A dehydrogenase; COX4, cytochrome C oxidase subunit 4. Error bars are 1 standard deviation (n = 10)

acute training stimulus in the form of greater mRNA levels of PGC-1 α and vascular endothelial growth factor.^{27,28} The insufficient muscular oxygen availability during exercise, causing local metabolic stress, is therefore probably highly significant in the oxidative and angiogenic improvements seen after endurance training. Lack of differences in training adaptation indicators between legs in the present study may therefore be explained by equal metabolic stress during training. This is supported by equal leg deoxy- and total hemoglobin signals in the present study, measured by near-infrared spectroscopy in four out of 11 subjects (unpublished data). This indicates preserved blood flow in both legs, which also other studies have confirmed with invasive measures.^{8,20}

However, in the present study it also may be questioned whether the difference in systemic variables between the 1L and 1L2A training exercises was sufficient to provide a differentiation in training response. This implies also that an increase in circulating catecholamines of 2-3 times more than that achieved during 1L is required for local training adaptations to take place. On the other hand, exercise protocols with even higher systemic response than during 1L2A would probably challenge oxygen delivery to the exercising leg due to blood flow restrictions, as first shown by Secher et al,²⁹ and thus threaten our purpose of achieving equal muscular metabolic stress in the legs. This makes the importance of differences in systemic variables per se on training adaptations difficult to investigate. However, in studies manipulating the amount of sympathetic response pharmacologically by administering β -adrenergic blockers, there are conflicting results in both acute responses to exercise and adaptations to training.⁹⁻¹⁵ Although the protocols in most cases are workload- or intensity-matched, there is no consensus, most likely because of differences in the degree of adrenergic blockade and dissimilarities in cardiovascular responses, together with the relative exercise capacities of subjects treated with β -blockers and control subjects.³⁰ It is also important to note that this study has some limitations. The lack of pre-training biopsies prevents investigation of muscular adaptations. In addition can the contralateral training protocol used not evaluate cross-transfer effects between the legs. The short break (10 minutes) between training modalities was enough to decrease catecholamine levels, but other factors, for example, different metabolites and cytokines, could still be upregulated and potentially interfere with training adaptations on the contralateral leg.

In conclusion, one-leg training with equal amounts of work and power output resulted in similar training responses in the legs regardless of whether arm cycling was added to the training or not. This indicates that local muscular training adaptations take place to the same extent in the presence of differences in systemic variables.

5 | PERSPECTIVES

How peripheral adaptations to endurance training are stimulated is still far from fully understood. Further studies are necessary to evaluate the different mandatory factors contribution and importance. Knowledge in this area is valuable when planning effective training protocols not only for elite athletes, but also for patient groups with different phenotypes and needs.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the staff of the Section of Specialized Endocrinology at Oslo University Hospital for their valuable cooperation. The authors have no conflicts of interest.

ORCID

Knut Sindre Mølmen  <https://orcid.org/0000-0001-8924-6848>

REFERENCES

1. Bassett DR, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc.* 2000;32(1):70-84.
2. Andersen P, Henriksson J. Capillary supply of the quadriceps femoris muscle of man: adaptive response to exercise. *J Physiol.* 1977;270(3):677-690.
3. Blomstrand E, Krstrup P, Søndergaard H, Rådegran G, Calbet J, Saltin B. Exercise training induces similar elevations in the activity of oxoglutarate dehydrogenase and peak oxygen uptake in the human quadriceps muscle. *Pflugers Arch Eur J Physiol.* 2011;462(2):257-265.
4. Desplanches D, Hoppeler H, Tüscher L, et al. Muscle tissue adaptations of high-altitude natives to training in chronic hypoxia or acute normoxia. *J Appl Physiol.* 1996;81(5):1946-1951.
5. Kiens B, Essen-Gustavsson B, Christensen NJ, Saltin B. Skeletal muscle substrate utilization during submaximal exercise in man: effect of endurance training. *J Physiol.* 1993;469:459-478.
6. Rådegran G, Blomstrand E, Saltin B. Peak muscle perfusion and oxygen uptake in humans: importance of precise estimates of muscle mass. *J Appl Physiol.* 1999;87(6):2375-2380.
7. Andersen P, Saltin B. Maximal perfusion of skeletal muscle in man. *J Physiol.* 1985;1985(366):233-249.
8. Savard GK, Richter EA, Strange S, Kiens B, Christensen NJ, Saltin B. Norepinephrine spillover from skeletal muscle during exercise in humans: role of muscle mass. *Am J Physiol.* 1989;257(6 Pt 2):H1812-H1818.
9. Miura S, Kawanaka K, Kai Y, et al. An increase in murine skeletal muscle peroxisome proliferator-activated receptor- γ co-activator-1 α (PGC-1 α) mRNA in response to exercise is mediated by β -adrenergic receptor activation. *Endocrinology.* 2007;148(7):3441-3448.
10. Robinson MM, Bell C, Peelor FF, Miller BF. β -Adrenergic receptor blockade blunts postexercise skeletal muscle mitochondrial protein

- synthesis rates in humans. *Am J Physiol Integr Comp Physiol*. 2011;301(36):327-334.
11. Ji LL, Lennon DL, Kochan RG, Nagle FJ, Lardy HA. Enzymatic adaptation to physical training under beta-blockade in the rat. Evidence of a beta 2-adrenergic mechanism in skeletal muscle. *J Clin Invest*. 1986;78(3):771-778.
 12. Svedenhag J, Henriksson J, Juhlin-Dannfelt A. Beta-adrenergic blockade and training in human subjects: effects on muscle metabolic capacity. *Am J Physiol*. 1984;247:E305-E311.
 13. Henriksson J, Svedenhag J, Richter EA, Christensen NJ, Galbo H. Skeletal muscle and hormonal adaptation to physical training in the rat: role of the sympatho-adrenal system. *Acta Physiol Scand*. 1985;123(2):127-138.
 14. Robinson MM, Richards JC, Hickey MS, et al. Acute β -adrenergic stimulation does not alter mitochondrial protein synthesis or markers of mitochondrial biogenesis in adult men. *AJP Regul Integr Comp Physiol*. 2010;298(1):R25-R33.
 15. Wolfel EE, Hiatt WR, Brammell HL, et al. Effects of selective and nonselective beta-adrenergic blockade on mechanisms of exercise conditioning. *Circulation*. 1986;74(4):664-674.
 16. Brandt N, Gunnarsson TP, Hostrup M, et al. Impact of adrenaline and metabolic stress on exercise-induced intracellular signaling and PGC-1 α mRNA response in human skeletal muscle. *Physiol Rep*. 2016;4(14):1-13.
 17. Fiorenza M, Gunnarsson TP, Hostrup M, et al. Metabolic stress-dependent regulation of the mitochondrial biogenic molecular response to high-intensity exercise in human skeletal muscle. *J Physiol*. 2018;596(14):2823-2840.
 18. Abbiss CR, Karagounis LG, Laursen PB, et al. Single-leg cycle training is superior to double-leg cycling in improving the oxidative potential and metabolic profile of trained skeletal muscle. *J Appl Physiol*. 2011;110(5):1248-1255.
 19. MacInnis MJ, Zacharewicz E, Martin BJ, et al. Superior mitochondrial adaptations in human skeletal muscle after interval compared to continuous single-leg cycling matched for total work. *J Physiol*. 2017;595(9):2955-2968.
 20. Richardson RS, Kennedy B, Knight DR, Wagner PD. High muscle blood flows are not attenuated by recruitment of additional muscle mass. *Am J Physiol*. 1995;269(5 Pt 2):H1545-H1552.
 21. Hallén J, Saltin B, Sejersted OM. K⁺ balance during exercise and role of beta-adrenergic stimulation. *Am J Physiol*. 1996;270(6 Pt 2):R1347-R1354.
 22. Andersen P, Adams RP, Sjøgaard G, Thorboe A, Saltin B. Dynamic knee extension as model for study of isolated exercising muscle in humans. *J Appl Physiol*. 1985;59(5):1647-1653.
 23. Ohnishi A, Minegishi A, Sasaki T, Suganuma T, Ishizaki T. Effect of beta-adrenoceptor blockade on exercise-induced plasma catecholamine concentrations and their dissipation profile. *Br J Clin Pharmacol*. 1987;23(3):339-343.
 24. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982;14(5):377-381. <http://www.ncbi.nlm.nih.gov/pubmed/7154893>.
 25. Paulsen G, Cumming KT, Holden G, et al. Vitamin C and E supplementation hampers cellular adaptation to endurance training in humans: a double-blind, randomised, controlled trial. *J Physiol*. 2014;592:1887-1901.
 26. Baar K. Nutrition and the adaptation to endurance training. *Sport Med*. 2014;44(Suppl. 1):S5-S12.
 27. Norrbom J, Sundberg CJ, Ameln H, Kraus WE, Jansson E, Gustafsson T. PGC-1 α mRNA expression is influenced by metabolic perturbation in exercising human skeletal muscle. *J Appl Physiol*. 2004;96(1):189-194.
 28. Gustafsson T, Ameln H, Fischer H, Sundberg CJ, Timmons JA, Jansson E. VEGF-A splice variants and related receptor expression in human skeletal muscle following submaximal exercise. *J Appl Physiol*. 2005;98(6):2137-2146.
 29. Secher NH, Clausen JP, Klausen K, Noer I, Trap-Jensen J. Central and regional circulatory effects of adding arm exercise to leg exercise. *Acta Physiol Scand*. 1977;100(3):288-297.
 30. Ihsan M, Watson G, Abbiss C. PGC-1 α Mediated Muscle Aerobic Adaptations to Exercise, Heat and Cold Exposure. *Cell Mol Exerc Physiol*. 2014;3(1):1-13.

How to cite this article: Mølmen KS, Hallén J, Rud B. Peripheral adaptations to endurance training—Effect of active muscle mass. *Transl Sports Med*. 2019;2:240–247. <https://doi.org/10.1002/tsm2.87>